ANNUAL REVIEWS

Annual Review of Earth and Planetary Sciences

Earth's Continental Lithosphere Through Time

Chris J. Hawkesworth,^{1,2} Peter A. Cawood,^{2,3} Bruno Dhuime,^{1,4} and Tony I.S. Kemp⁵

¹School of Earth Sciences, University of Bristol, Bristol BS8 1RJ, United Kingdom; email: c.j.hawkesworth@bristol.ac.uk, B.Dhuime@bristol.ac.uk

²School of Earth and Environmental Sciences, University of St. Andrews, St. Andrews KY16 9AL, United Kingdom

³School of Earth, Atmosphere & Environment, Monash University, Melbourne, VIC 3800, Australia; email: peter.cawood@monash.edu

⁴CNRS-UMR 5243, Géosciences Montpellier, Université de Montpellier, 34095 Montpellier Cedex 05, France

⁵School of Earth Sciences, University of Western Australia, Perth, WA 6009, Australia; email: tony.kemp@uwa.edu.au

Annu. Rev. Earth Planet. Sci. 2017. 45:169-98

First published as a Review in Advance on May 31, 2017

The Annual Review of Earth and Planetary Sciences is online at earth.annualreviews.org

https://doi.org/10.1146/annurev-earth-063016-020525

Copyright © 2017 by Annual Reviews. All rights reserved



ANNUAL **Further** Click here to view this article's online features:

- Download figures as PPT slides
 Navigate linked references
- Navigate linked referer
 Download citations
- Download citations
 Explore related articles
- Search keywords

Keywords

continental crust, continental mantle lithosphere, plate tectonics, preservation bias, early Earth

Abstract

The record of the continental lithosphere is patchy and incomplete; no known rock is older than 4.02 Ga, and less than 5% of the rocks preserved are older than 3 Ga. In addition, there is no recognizable mantle lithosphere from before 3 Ga. We infer that there was lithosphere before 3 Ga and that \sim 3 Ga marks the stabilization of blocks of continental lithosphere that have since survived. This was linked to plate tectonics emerging as the dominant tectonic regime in response to thermal cooling, the development of a more rigid lithosphere, and the recycling of water, which may in turn have facilitated plate tectonics. A number of models, using different approaches, suggest that at 3 Ga the volume of continental crust was \sim 70% of its presentday volume and that this may be a minimum value. The continental crust before 3 Ga was on average more mafic than that generated subsequently, and this pre-3 Ga mafic new crust had fractionated Lu/Hf and Sm/Nd ratios as inferred for the sources of tonalite-trondhjemite-granodiorite and later granites. The more intermediate composition of new crust generated since 3 Ga is indicated by its higher Rb/Sr ratios. This change in composition was associated with an increase in crustal thickness, which resulted in more emergent crust available for weathering and erosion. This in turn led to an increase in the Sr isotope ratios of seawater and in the drawdown of CO_2 . Since 3 Ga, the preserved record of the continental crust is marked

by global cycles of peaks and troughs of U-Pb crystallization ages, with the peaks of ages appearing to match periods of supercontinent assembly. There is increasing evidence that the peaks of ages represent enhanced preservation of magmatic rocks in periods leading up to and including continental collision in the assembly of supercontinents. These are times of increased crustal growth because more of the crust that is generated is retained within the crust. The rates of generation of continental crust and mantle lithosphere may have remained relatively constant at least since 3 Ga, yet the rates of destruction of continental crust have changed with time. Only relatively small volumes of rock are preserved from before 3 Ga, and so it remains difficult to establish which of these are representative of global processes and the extent to which the rock record before 3 Ga is distorted by particular biases.

1. INTRODUCTION

The lithosphere is the rigid outer layer of Earth, and it is divisible into crust and upper mantle components (Figure 1). The formation and evolution of the lithosphere are driven by thermal energy from the planet's interior and modulated by solar radiation, which regulates the temperature at the surface of the lithosphere. The lithosphere records interactions with the underlying asthenospheric mantle and with the overlying hydrosphere and atmosphere. The lithosphere is the long-term archive of Earth's history, and yet our knowledge of this history is limited by variable but generally poor preservation and by difficulties in accessing its deep crustal and mantle components. The rock record is incomplete, and the lithosphere that is preserved may not be representative of the processes that formed it (Hawkesworth et al. 2016). The rocks that are readily accessible for sampling record long and complex cycles of generation and reworking, with most crustal rocks derived from preexisting crustal rocks, from "the ruins of mountains" (Hutton 1788, p. 215). Hence, it remains difficult to see back through to the composition and to the tectonic settings in which the continental lithosphere was generated. In this contribution, we consider the nature of the continental crust and its mantle lithosphere, focusing on the former, for which there are more data, and then discuss the interrelationship between the two and the tectonic processes that have controlled their evolution.

At the present, the distinction between the continental and oceanic lithosphere is marked. The oceanic lithosphere is relatively dense, and its crustal component is mafic and less than 200 Myr old (Cloos 1993, Kearey et al. 2009). In contrast, the continental lithosphere is more differentiated and buoyant and is harder to destroy; the crust preserves rocks that are up to 4.02 Ga and zircons that are up to 4.4 Ga in age (Bowring & Williams 1999, Reimink et al. 2016, Wilde et al. 2001). Half of the exposed rocks in the continental crust have formation ages (e.g., igneous crystallization ages for magmatic rocks, or deposition ages for sedimentary rocks, referred to here as geological ages) less than 600 Ma, and fewer than 5% have geological ages older than 3.0 Ga (Goodwin 1996). Before 3.0 Ga, more of Earth's crust appears to have been mafic, and the distinction between continental crust older than 2.7 Ga. The continental crust evolved from a predominantly mafic composition to today, when it is andesitic and consists of three generalized layers: the upper, middle, and lower crust (Dhuime et al. 2015, Rudnick & Gao 2003, Tang et al. 2016). See Hacker et al. (2015) for an alternative two-layer crust.

Figure 2 summarizes the ages (circles) and age ranges (bars) of major events and cycles in Earth history. It highlights the nature of the geological record through the peaks of zircon crystallization



Schematic cross section of types of continental lithosphere emphasizing the thick stable nature of Archean cratons, accompanied by a glossary of some of the terms used. Thickness of lithosphere beneath Archean regions is of the order of 200–250 km, and that of the oceanic lithosphere is up to 100 km (after Cawood et al. 2013).

ages and the distribution of mantle depletion ages, the cyclical pattern of supercontinent formation, and the apparently intermittent nature of Au and volcanic massive sulfide mineralization. The early crust is not recorded, but it is assumed to have developed from the initial magma ocean and to have been mafic in composition (e.g., Elkins-Tanton 2008). The oldest zircon at 4.4 Ga indicates that there were felsic magmas at that time, and the elevated δ^{18} O values in some zircon crystals are taken to be evidence for the presence of surface water by 4.4 Ga (Valley et al. 2002, Wilde et al. 2001). Mafic and ultramafic volcanic rocks in greenstone belts are associated with tonalite-trondhjemite-granodiorite rocks in the classic granite-greenstone belt terrains of the Archean. The late Archean was characterized by tonalite-trondhjemite-granodiorite magmatism, the remelting of intermediate to felsic crust, the widespread generation of relatively potassic granites, and the stabilization of the continental crust and mantle lithosphere (Anhaeusser & Robb 1981, Carlson et al. 2005, Dewey & Windley 1981, Schoene et al. 2009).

The lithosphere is described on the basis of its mechanical properties as Earth's rigid outer layer, which can support horizontal and vertical stresses. It is recognized as a thermal boundary layer that cools by conduction, as a seismic lithosphere marked by an abrupt decrease in shear wave velocity at its base (a low-velocity zone), and as an elastic layer whose thickness is determined from the response to surface loads, such as seamounts and mountain belts (Watts et al. 2013).

Geotherms are steeper in the continents, and so the thermal boundary layer is thicker beneath continents than beneath the oceans. The continental mantle lithosphere shows significant variations as surface heat flux values increase, and thickness decreases, with decreasing lithospheric age. Present-day heat flux values range from $30-40 \text{ mW/m}^2$ for Archean regions with lithosphere thicknesses of >150 km to >60-80 mW/m² and generally <100 km for Phanerozoic regions (Jaupart & Mareschal 2003, Jaupart et al. 1998, McLennan et al. 2005, Nyblade 1999, Pollack et al. 1993). Moreover, global versus seismic tomography models (Grand 2002, Shapiro & Ritzwoller 2002) and the global thermal model of Artemieva & Mooney (2001) yield consistent estimates of the lithospheric thickness beneath continents (Artemieva 2009).



The continental lithospheric mantle beneath Archean-aged crust tends to be composed of dehydrated, highly depleted mantle peridotite (Boyd 1989, Boyd et al. 1997, Griffin et al. 2009, Pearson et al. 1995, Pollack 1986), resulting in it being intrinsically buoyant and strong, which counteracts the destabilizing effect of its cold thermal state (Jordan 1978, 1988). These regions of thick ($> \sim 150$ km) stable lithosphere are referred to as cratons (kratogen) (**Figure 1**) (Kober 1921), and owing to their association with Archean crust, as well as Archean ages from xenoliths of underlying mantle lithosphere (Pearson & Wittig 2013), they are largely thought to have formed at this time. From a geological perspective, cratons may be viewed as areas of stable continental lithosphere in which weakly deformed and little metamorphosed Proterozoic sedimentary rocks unconformably overlie a deformed and metamorphosed Archean basement (e.g., Moores & Twiss 1995). On the basis of recent interpretations of Rayleigh wave tomography data, McKenzie et al. (2015) show that thick lithosphere (>200 km) remains a feature of Archean terrains and that areas of 150–200-km thick lithosphere are present under the Proterozoic crust. McKenzie & Priestley (2008) refer to the thick lithosphere currently developed beneath Tibet as a proto-craton, implying the process of craton formation is in part related to continental collision and continues today.

2. OROGENIC AND GLOBAL CYCLES

Geological cycles operate on a range of spatial and temporal scales. James Hutton (1785, 1788) concluded that there had been innumerable cycles, involving deposition on the seabed, uplift with tilting and erosion, and submergence under water for additional layers to be deposited. Orogenic cycles provide a framework whereby periods of sediment deposition are followed by compression, crustal thickening, uplift, and erosion. They differ in spatial and temporal scales from supercontinent cycles, which are global features and are therefore more likely to have had a significant role in the evolution of the continental crust as a whole. Supercontinent cycles have punctuated the post-Archean history of Earth, and they involve the episodic assembly and breakup of large landmasses, with profound consequences for the geologic record of both the solid and surficial Earth (Nance et al. 2014, Worsley et al. 1986).

Zircon remains the key timepiece for dating events in the geological record, and hundreds of thousands of U-Pb zircon ages have been published (e.g., Voice et al. 2011). Zircon typically crystallizes from relatively felsic magmas, and it is preserved in detrital sediments that have sampled continental crust. Compilations of age data have long been known to yield episodic

Figure 2

Ages (*circles*) and age ranges (*bars*) of major events and cycles preserved in Earth history. Sources of the data are indicated in the figure key: (1) The oldest terrestrial fragments, zircons from the Jack Hills; (2) the age range of principal peaks in U-Pb crystallization ages; (3) the oldest rocks, Slave craton; (4) the approximate age range of supercontinent assembly; (5) the late heavy bombardment; (6) the age range of eclogitic inclusions in diamonds; (7) the oldest Os model ages from detrital osmiridium grains, Witwatersrand Basin; (8) Re-depletion age ranges for peridotite xenoliths (see also **Figure 5**); (9) the \sim 3.8 Ga Nuvvuagittuq greenstone belt, Superior craton, and 3.85 Ga metabasalts from Isua, Greenland, are similar to modern subduction-related islands; (10) the pre-3.2 Ga rock units from Pilbara craton are associated with vertical tectonics, and rocks younger than 3.1 Ga are inferred to have formed by plate subduction processes; (11) the shift from predominantly mafic crustal compositions prior to 3.0 Ga to increasingly felsic compositions by 2.5 Ga; (12) the secular increase in time-integrated Rb/Sr ratios between 3.1 Ga and 1.7 Ga, indicative of a doubling in average continental crustal thickness from ca. 20 km to ca. 40 km; (13) the main pulses of orogenic gold deposition; (14) the main pulses of volcanic-hosted massive sulfide deposits; (15) subaerial large igneous province magmatism increases from 3.0 to 2.5 Ga, corresponding with the increase in 87 Sr/⁸⁶Sr ratios of seawater consistent with the emergence and weathering of continental crust; (16) the significant difference between the relative paleopositions of the Kaapvaal and Superior cratons at 2.68 Ga and 2.07 Ga and Baltica and Australia between 1.77 Ga and 1.5 Ga; (17) the high-pressure and ultrahigh-pressure metamorphism, which are limited to rock units dated at <0.7 Ga; and (18) the increases in atmospheric oxygen values during the Great Oxidation Event and the Neoproterozoic Oxidation Event.

distributions (Gastil 1960, Stockwell 1961), but these were based on mineral systems that can be reset through younger tectonothermal events (e.g., K-Ar, Ar-Ar, Rb-Sr). However, this episodic distribution is also a characteristic of global compilations of physiochemical resistant zircon U-Pb ages (**Figure 3**). Such apparently cyclical patterns are unexpected in a plate tectonic Earth in which broadly similar volumes of magmas are likely to have been generated, and destroyed, each year (Scholl & von Huene 2009).

The peaks and troughs of zircon ages have been interpreted as primary signals of periods of high and low volumes of granitic magmas, perhaps related to mantle superplumes (Albarède 1998, Arndt & Davaille 2013, Condie 1998, Condie et al. 2015, Parman 2015, Rino et al. 2004). There are, however, a number of observations that question this interpretation. First, the distribution of ages is based on a global data set, and so it would appear necessary to invoke periods of increased plume activity on a large scale. Second, there are peaks of zircon crystallization ages at ~1 Ga, 0.6 Ga, and 0.3 Ga (**Figure 3**), corresponding with the Grenville, Pan-African, and Variscan-Alleghenian orogenies, which are widely regarded as periods of continental assembly and collision (e.g., Cawood & Buchan 2007, Cawood et al. 2016, Gower & Krogh 2002), rather than times of unusual volumes of magma generation. Third, the composition of the continental crust is dominated by minor and trace element features that are characteristic of subduction-related magmas rather than those generated in intraplate settings (Rudnick & Gao 2003).

Large volumes of magma are generated above subduction zones, but in these settings, continental crust is also destroyed by erosion, subduction, and in some areas density foundering at rates similar to, or greater than, those at which new crust is generated (Clift et al. 2009; Scholl & von Huene 2007, 2009; Stern 2011). The existence and ages of different supercontinents are still matters for debate (e.g., Arndt & Davaille 2013, Bradley 2011, Campbell & Allen 2008), but there is a broad link between the peaks of U-Pb zircon ages, periods of crustal reworking as determined from zircon Hf and O isotopic records, and the inferred development of supercontinents (**Figure 3**). We therefore prefer models in which the peaks of zircon ages reflect the preferential preservation of magmatic rocks formed during the latter stages of ocean closure that mark the times of supercontinent assembly (Cawood et al. 2013, Hawkesworth et al. 2009). These would then be times in which the volume of stabilized continental crust would be expected to increase not because more new crust had been generated but because more of the crust that had been generated was retained within the continental crust. Overall the supercontinent cycle inherently biases the rock record both through selective isolation of material in continental cores during supercontinent assembly and through removal and recycling of material formed during stages of extension and convergence.

The assembly of supercontinents involves continental collision, and the associated thickening of the continental crust should result in increased amounts of crustal reworking. One proxy for

Figure 3

(*a*) Histogram of over 100,000 detrital zircon analyses showing peaks in U-Pb crystallization ages over the course of Earth history (Voice et al. 2011), which are similar to the ages of supercontinent assembly. Also shown is the apparent thermal gradient at the age of peak metamorphism for the three main types of extreme metamorphic belts (Brown 2007, 2014). (*b*) Normalized seawater 87 Sr/ 86 Sr curve (Shields 2007), and the running mean of initial ε Hf in ~7,000 detrital zircons from recent sediments (Cawood et al. 2013). (*c*) The red curve represents changes in the degrees of reworking of the continental crust through time, calculated from the distributions of the proportions of reworked crust and new crust, calculated from Hf isotope variations in zircons (Dhuime et al. 2012). The blue curve is the moving average distilled from compilation of ~3,300 δ^{18} O analyses of zircon versus U-Pb ages (Spencer et al. 2014). Abbreviations: HP, high-pressure; UHP, ultrahigh-pressure; UHT, ultrahigh-temperature. ε Hf is a measure of the deviation of the Hf isotope ratios of a sample from that of a chondritic reservoir, taken to be the bulk earth, as expressed in

$$\varepsilon \text{Hf} = \left[\frac{\binom{1/6}{\text{Hf}} \frac{1}{1/1} \frac{1}{\text{Hf}} \frac{1}{\text{Sample}} - \binom{1}{6} \frac{1}{\text{Hf}} \frac{1}{1/1} \frac{1}{\text{Hf}} \frac{1}{\text{Chon}}}{\binom{1}{6} \frac{1}{\text{Hf}} \frac{1}{1} \frac{1}{1} \frac{1}{\text{Hf}} \frac{1}{1} \frac{1}{1$$



crustal reworking is the temporal distribution of crystallization ages of zircons with Hf model ages greater than their crystallization ages (Dhuime et al. 2012). Another is the distribution of the peaks in δ^{18} O values in zircons through time (**Figure 3**) (Bindeman et al. 2016, Dhuime et al. 2012, Roberts & Spencer 2014, Spencer et al. 2014). Elevated δ^{18} O values indicate reworking of supracrustal material, and this is most readily achieved in sections of crust thickened in response to continental collision. Both proxies indicate that the periods of increased crustal reworking are also those of supercontinent assembly (**Figure 3**). Thus, this is independent evidence that the peaks of U-Pb crystallization ages are associated with periods of crustal thickening and of continental collision, and the development of supercontinents.

3. SAMPLING THE CONTINENTAL CRUST

Zircons provide a record of the ages of rocks exposed on Earth's surface, and there is reasonable agreement on the composition of the upper and the bulk of the continental crust (Rudnick & Gao 2003; Taylor & McLennan 1985, 1995). The crust can be sampled by analyzing clastic sediments derived by weathering and erosion of different rock units in their source areas. The compositions of clastic sedimentary rocks provide reasonable estimates of the average minor and trace element contents of their source rocks, at least for less soluble elements (McLennan et al. 2005, Taylor & McLennan 1985). However for radiogenic isotopes, the ages of the different source rocks become important, and it is necessary to establish the extent to which source rocks with different ages contribute to the sediments analyzed. Specifically, it is necessary to determine whether the proportions of source rocks in the eroded terrain are similar to those estimated from bulk analysis of the resultant sediment (Allègre & Rousseau 1984). If not, it becomes critical to correct for the bias introduced during sedimentary processes to unravel the age profile of the upper continental crust.

Archean rocks within the drainage of the modern-day Frankland River in southwest Australia are, for example, underrepresented by a factor of approximately six in the sediments sampled at the river mouth (Cawood et al. 2003, Dhuime et al. 2011). The post-Archean rocks are exposed in areas of relatively high relief, and where the river contains larger volumes of water, and so they are likely to contribute disproportionately large volumes of detritus to the Frankland River, with the result that the Archean crust is underrepresented. **Figure 4** shows that, in general, crust with Archean Nd model ages tends to have lower average topographic relief and that the average Nd model age of samples taken from areas with more than 100 m of relief is ~ 1 Ga (Allan 2014). Given that denudation rates increase with topographic relief (Larsen et al. 2014) and that the average model age of at least the upper continental crust is thought to be ~ 1.8 Ga (Chauvel et al. 2014), it follows that old rocks are significantly underrepresented in the sedimentary record.

Allègre & Rousseau (1984) addressed the issue of bias introduced by using detrital sediments to sample the continental crust. They defined an erosion parameter K to relate the model age of the sediments analyzed to the average model age of their source rocks. For a source terrain with two units of different model ages, K is the ratio of the material from the younger to the older terrain determined from analyses of the sediment divided by the ratio of the area of the younger relative to the older terrain present in the catchment area, i.e., estimated from a geological map. This has proven difficult to measure in natural systems, and values ranging between two and three have been assumed in previous studies (Allègre & Rousseau 1984, Garrels & Mackenzie 1971, Goldstein & Jacobsen 1988, Jacobsen 1988, Kramers 2002, Kramers & Tolstikhin 1997). Such values for K would suggest that the volume of continental crust at 3 Ga was ~25–30% of the present volume. However, using the value of K = 15, obtained from the sediments at the mouth



Present-day Nd isotope ratios for whole-rock samples plotted against topographic relief (Allan 2014). The average Nd model age for samples from areas of >100-m relief is ~ 1 Ga, whereas estimates of the average Nd model age for the continental crust is ~ 1.8 Ga from Condie & Aster (2010) and Chauvel et al. (2014).

of the Frankland River (Dhuime et al. 2011), suggests that the volume of continental crust present at 3 Ga was 65–70% of the present volume.

4. MANTLE LITHOSPHERE AND ITS RELATIONSHIP TO THE OVERLYING CRUST

The mantle lithosphere is the rigid portion of mantle attached to the overlying crust (**Figure 1**), and so it may preserve a record of upper mantle processes dating back to the formation of the oldest preserved continental crust. It can be mapped by geophysics (e.g., James et al. 2001, Lee & Nolet 1997, McKenzie et al. 2015, McKenzie & Priestley 2008, Mooney et al. 1998), and it is sampled at Earth's surface as fragments transported in younger kimberlites and alkali basalts (e.g., Carlson et al. 2005). Mineral thermobarometry on suites of samples of mantle lithosphere from adjacent localities has been used to reconstruct geotherms within the lithosphere. Typically, the lowest geotherms are found beneath stable cratons where temperatures at 100 km depth are between 750°C and 900°C (Carlson et al. 2005, Finnerty & Boyd 1987, Lee et al. 2011).

Many mantle peridotites have undergone alteration and/or metasomatism, and the Mg number of olivine is taken as the most reliable indication of the melting processes responsible for their initial bulk rock compositions (Pearson & Wittig 2013). The variation of the Mg number of olivine with depth in the lithosphere can be compared with experimental parameterizations of mantlemelting products under different conditions. Peridotite xenoliths from Archean crustal terrains tend to have olivines with higher Mg numbers, and hence, these are more residual in composition than those from massif peridotites, which are tectonically exhumed fragments of the upper mantle (Bodinier & Godard 2003), and those mantle xenoliths from off-craton localities. The xenoliths from Archean areas tend to have lower Fe contents, and hence, these are less dense and therefore more buoyant than those in younger areas. Many of these xenoliths also have elevated silica (Boyd 1989 and references therein, Carlson et al. 2005, Pearson & Wittig 2013). The Archean lithosphere is thicker than younger counterparts; it is less dense and hence harder to destroy than the more Fe-rich mantle, and so it may have had a role in the long-term survival of the Archean continental crust (Davidson & Arculus 2006).

The generation of low-Fe (high-Mg number) residual mantle in the Archean has been the subject of much debate. Walter (1998) demonstrated experimentally that a low-Fe residual mantle can result from the generation of komatiite magmas from fertile peridotite at depths >7 GPa (~250 km). However, Pearson & Wittig (2013) highlighted that the low heavy rare earth element (REE) content, and the fractionation of Lu from Yb, in most mantle xenoliths from cratonic areas can only be generated by extensive melting in the spinel peridotite stability field even though garnet is present as a residual phase in Walter's experiments. Herzberg & Rudnick (2012) invoked a model of non-arc magmatism with mantle potential temperatures of up to $\sim 1,700^{\circ}$ C, and melt fractions of 0.25–0.45, at the time of formation of cratonic mantle lithosphere. Variations in olivine Mg number with depth are consistent with models in which the peridotites represent residues after melting at high temperatures at shallow depths, as in a hot ridge setting, and there is relatively little evidence for the even higher Mg number residues than would be expected in a hot mantle plume setting (Herzberg & Rudnick 2012, Pearson & Wittig 2013). Intriguingly in the context of when plate tectonics might have developed, Simon et al. (2007) developed a model for the origins of low-temperature peridotite xenoliths from Kimberley and Lesotho involving shallow ridge melting, followed by subduction-related metasomatism probably during amalgamation of smaller preexisting terranes in the Late Archaean (~ 2.9 Ga).

Matsukage & Kawasaki (2014) argued that cratonic garnet peridotites could reflect melt depletion from almost anhydrous to a maximum of approximately 1% to 2% water content in the upper mantle at depths of 100 to 200 km. Such elevated water content would imply the presence of subduction, and eclogites are fragments of basalt, which are also interpreted as evidence for subduction. Eclogites have been dated back to 2.9 Ga (Tappe et al. 2011), and many have Os model ages of 2.7–3.0 Ga. Eclogite assemblages are recognized in inclusions in diamonds back to 3 Ga and not before (Shirey & Richardson 2011). Eclogite provides evidence of the return of basalt to the mantle, and although this is commonly attributed to the onset of subduction (Shirey & Richardson 2011), it might also happen through density foundering of crustal material.

Melt depletion events in mantle rocks may now be dated using the Re-Os decay scheme. Walker et al. (1989) defined the Re-depletion model age, T_{RD} , calculated by assuming that the Re/Os ratio of the analyzed samples was zero since the time of melt extraction. This in effect provides a minimum age for the depletion event, as not all the Re may have been removed.

Mantle xenoliths from cratonic areas tend to have older T_{RD} ages, with a peak at 2.8 Ga, and off-craton xenoliths have younger T_{RD} ages with peaks at 2.4 and 1.5 Ga. Spinel peridotites tend to occur in relatively young geological settings, typically in alkali basalts rather than kimberlites, and most T_{RD} ages from spinel peridotites are less than 1.5 Ga (**Figure 5**). It is tempting to assume that mantle depletion events reflect the time of formation of the continental mantle lithosphere, but that is not always the case as some segments of mantle within the asthenosphere also have depleted Os isotope signatures before they are incorporated into the lithosphere (Liu et al. 2015). Nonetheless, as noted by many studies (Carlson et al. 2005, Pearson & Wittig 2013, and references



Histogram of Re-depletion (T_{RD}) ages from on and off craton and from spinel peridotites. The data for on craton peridotites are from Pearson & Wittig (2013 and the references therein), and the other sources of data are as follows: Becker et al. (2001, 2006); Burton et al. (2000); Carlson & Irving (1994); Carlson et al. (2004); Chesley et al. (1999); Choi et al. (2010); Gao et al. (2002); Handler et al. (1997, 2003, 2005); Janney et al. (2010); Lee et al. (2000); Liu et al. (2011, 2015, 2016); Luguet et al. (2009); Meisel et al. (2001); Pearson & Wittig (2013); Peslier et al. (2000a,b); Simon et al. (2007); Wu et al. (2003, 2006); Xu et al. (2008); H.-F. Zhang et al. (2009); and Y.-L. Zhang et al. (2011).

therein), mantle rocks with Archean T_{RD} ages predominate in areas of Archean crust, and those with younger T_{RD} ages generally occur beneath younger continental crust.

5. CONSTRAINING THE VOLUME OF CONTINENTAL CRUST AND HOW THIS HAS CHANGED WITH TIME

The issue of the volume of continental crust through time has been a long-standing debate (Armstrong 1981, Gastil 1960, Moorbath 1977, O'Nions et al. 1979). Models of the progressive accretion of mobile belts onto older cratonic cores may have implied linear crustal growth through time (e.g., Dana 1873), and although early compilations of geochronological data appeared to support this concept (e.g., Hurley et al. 1962), in practice they reflected reworking of older crust by younger orogenic events. Other models were based on arguments that Earth has progressively cooled, and so a relatively large volume of crust was generated early and was subsequently destroyed and generated at rates similar to those observed today (Armstrong 1981). However, little of that old crust is preserved (**Figure 6**), and so the debate became linked to the



Mean curves from the continental growth models of Belousova et al. (2010), Dhuime et al. (2012), and Pujol et al. (2013) are compared with the distribution of geological ages in presently preserved crust from Goodwin (1996). The Condie & Aster (2010) curve reflects the proportion of juvenile crust of different ages preserved today. The Belousova et al. (2010) and Dhuime et al. (2012) curves are based on the proportions of reworked and juvenile crust in the zircon record. The K = 1 curve is the continental growth curve for the Gondwana supercontinent calculated from the Nd isotope data for Australian shales (Allègre & Rousseau 1984), assuming an erosion parameter K = 1 (i.e., no preferential erosion of the different lithologies producing the sediment). A K value of unity would imply that 30% of the continental crust was generated by the end of the Archean, but this increases to 75% if K = 15 (curve from Cawood et al. 2013). The analyses of detrital zircons (Belousova et al. 2010, Dhuime et al. 2012) are not sensitive to values of K. K influences the proportions of old and young zircons available for analysis, but the crustal growth curves of Belousova et al. (2010) and Dhuime et al. (2012) are based on the proportions of reworked and juvenile crust irrespective of the age distribution of the zircons analyzed.

ease with which continental crust might be destroyed. The advent of Nd and then Hf isotope measurements allowed the age of formation of different segments of the continental crust to be investigated (Allègre & Rousseau 1984, Belousova et al. 2010, Dhuime et al. 2012, McCulloch & Wasserburg 1978, O'Nions et al. 1983). Yet many of the early Nd isotope studies relied on data from shales, and as discussed above in Section 3, older geological terrains are significantly underrepresented in the compositions of terrigenous sediments (**Figure 6**).

The distributions of crust formation ages (**Figure 3***a*) and of the proportions of reworked crust and how that has changed with time (**Figure 3***c*) do not indicate the volume of crust from different periods. Initial models regarded the continental crust as the complementary reservoir to the depleted upper mantle, and investigators sought to ensure that the volume of crust generated at different times would result in a residual depleted mantle reservoir with what was regarded as a reasonable average age of 2.0-1.8 Ga (Allègre et al. 1983, Jacobsen & Wasserburg 1979, Kramers & Tolstikhin 1997, O'Nions et al. 1979). Recent models have focused on the records available from the crust itself; they are anchored by the present-day volume of the crust and then projected back in time based on estimated ratios of the proportions of new and reworked crust for different time slices until the inferred time of initial formation of the continental crust (Belousova et al. 2010, Dhuime et al. 2012).

The proportions of new and reworked crust can be estimated from (a) the balance between zircons with Hf model ages close to their crystallization ages and those with model ages significantly older than their crystallization ages (Belousova et al. 2010, Dhuime et al. 2012), and from (b) Nd isotope ratios in clastic sediments (Allègre & Rousseau 1984, Dhuime et al. 2011). Approaches based on Nd and Hf model ages involve a number of uncertainties linked to how well the formation age of new continental crust is constrained by the calculation of model ages (Arndt & Goldstein 1987). There are also concerns that for detrital zircons of Archean age it remains difficult to establish whether they have suffered Pb loss, which would result in shifts to younger ages and thence spuriously old model ages (Vervoort & Kemp 2016). Nonetheless, two independent approaches, one based on Hf isotopes in zircons and one based on variations in the Nd isotope ratios of shales, yield similar curves for the changes in the volumes of continental crust through time (the curves labeled Dhuime et al. 2012 and K = 15 in Figure 6). The distinction is that the approaches of Belousova et al. (2010) and Dhuime et al. (2012) are concerned with the volumes of new crust that were present at different times in the past, whereas the Allègre & Rousseau (1984) model evaluates the proportion of crust with different model ages at the present day and infers that it is a fair reflection of the volumes of new crust that were present at different times in the past. Campbell (2003) used secular changes in Nb/U in mafic rocks as a nonisotope approach to constraining crustal volumes and concluded that by ~ 3 Ga the volume of continental crust was $\sim 70\%$ of the present volume of the continental crust. Pujol et al. (2013) reached a similar conclusion based on estimates of Archean atmospheric 40Ar/36Ar from measurements of Ar in 3.5-Ga hydrothermal quartz. Such values of \sim 70% of the crust at 3 Ga are significantly higher than the volume of such crust preserved at the present day [<5% from the Condie & Aster (2010) curve in Figure 6], which in turn highlights the question of how and when such volumes of old crust were destroyed (see Section 6.3).

6. SYNTHESIS

Data from field studies along with paleomagnetic, geochemical, and ore deposit studies indicate a change in the character, composition, and thickness of the continental lithosphere in the latter half of the Archean from \sim 3.2 Ga (**Figure 2**) (Cawood et al. 2006, Dhuime et al. 2012, Keller & Schoene 2012, Tang et al. 2016). There are no significant peaks of zircon crystallization ages older than 3 Ga (**Figure 3**), and the development of potassic granites from 3.1 to 2.6 Ga and main peak of old T_{RD} ages (3.0–2.7 Ga) (**Figure 5**) are considered to mark the stabilization of the continental lithosphere (Anhaeusser & Robb 1981, Carlson et al. 2005, Schoene et al. 2009). Recent curves for the volume of continental crust through time (**Figure 6**) are also marked by a change in slope at approximately 3 Ga, reflecting a change in the estimated growth rate of the continental crust. Such changes presumably happened over a period of time and at different times in different places. Nonetheless, 3 Ga is a useful time marker, and here, we discuss some of the differences that characterize geological terrains older and younger than 3 Ga.

6.1. Before 3 Ga

The oldest terrestrial zircons, from the Jack Hills area in Western Australia, indicate that there were felsic magmas and surface water from \sim 4.4 Ga (Valley et al. 2002, Wilde et al. 2001). There is, however, much controversy over the setting in which these Hadean zircons were generated and thus their significance for continental crustal growth (e.g., Harrison 2009, Kemp et al. 2010).

The oldest known terrestrial rocks, the Acasta gneisses, have components dated at \sim 4.02 Ga (Bowring & Williams 1999). At least in some Archean areas, pre-3.2 Ga crust is dominated by dome and basin granite-greenstone terranes, thought to be indicative of vertical tectonics (Bédard

et al. 2013, Van Kranendonk et al. 2007). Willbold et al. (2009) and Reimink et al. (2014) noted that Archean tonalitic gneisses are similar in composition to intermediate lavas generated in Iceland at the present day and invoked similar models linked to mantle upwelling. Pre-3.1 Ga upper crust also appears to have been more mafic than that generated subsequently (Tang et al. 2016, Taylor & McLennan 1985). Taylor & McLennan (1985) reported a sharp increase in Th/Sc in fine-grained sedimentary rocks at the end of the Archean. Tang et al. (2016) demonstrated that Archean shales and glacial diamictites have strikingly high Ni/Co and Cr/Zr ratios compared to younger sedimentary rocks. Both ratios correlate with MgO contents in igneous rocks, and Tang et al. (2016) concluded that the average upper crust, as sampled by these sediments, had ~15% MgO at 3.2 Ga and that this had dropped to ~4% by 2.6 Ga.

Geochemical studies on more mafic lithologies suggest that many of the older rock suites from Barberton, Pilbara, and West Greenland have affinity with modern within-plate magmas (Campbell 2003, Jenner et al. 2013, Næraa et al. 2012, Smithies et al. 2007, Van Kranendonk et al. 2007). Old mafic rocks with chemical similarities to recent subduction-related suites have also been reported, and at least for the period 3.85-3.2 Ga, they appear confined to West Greenland and Canada (Jenner et al. 2009, O'Neil et al. 2011). Thus, the magmatic records appear different in different places. In southern Africa and the Pilbara, there appears to be a shift from early within-plate-like magmas and an emphasis on vertical tectonics (Campbell 2003, Sandiford et al. 2004, Smithies et al. 2007, Van Kranendonk et al. 2007) to magmatic rocks with subductionrelated signatures after \sim 3.1 Ga (Smithies et al. 2007). In the Superior province, the older rocks have subduction-related affinities, and in West Greenland there are >3.8 Ga rocks with both subduction-related and within-plate geochemical signatures (Jenner et al. 2009, 2013). At present it appears that Archean areas with the classic dome and basin structures, as in the East Pilbara and southern Africa, are more likely to have within-plate magmatic rocks, associated with dominantly vertical tectonic regimes (e.g., Sandiford et al. 2004).

Very few rocks, and even crystals of zircon, have survived from before \sim 3.9 Ga, a time that marks the end of the period of heavy meteorite bombardment (**Figure 2**). Yet Earth would have had a crust before this, and it is likely that it would have been thick enough to have remelted and so developed a felsic component (Kamber 2015, Kemp et al. 2010). Marchi et al. (2014) rescaled the lunar meteorite flux to Earth and argued that the surface of the Hadean Earth was widely reprocessed by impacts through mixing and burial by impact-generated melt. Hansen (2015) took the argument further and contrasted Archean granite-greenstone terrains on Earth with crustal plateaus now recognized on Venus. In Hansen's (2015) model, Archean cratons are formed in response to a large bolide-piercing thin Archean lithosphere, causing localized high-temperature, high-fraction partial melting in the sublithospheric mantle and the melt rises, forming an igneous province that evolves to form a granite-greenstone terrain, whereas the melt residue develops a complementary mantle lithosphere.

Such models remain difficult to test, but the available evidence suggests that there was both subduction-related and what is typically termed within-plate magmatism at similar times in different terranes in the early Archean, before subduction-related magmatism became more dominant after \sim 3.2 Ga.

6.2. Changes About 3 Ga

The geological record is characterized by peaks and troughs in the distribution of U-Pb zircon crystallization ages from 2.7 Ga to the present (**Figure 3**). The peaks of ages are taken to reflect periods of continental collision and crustal thickening in the development of supercontinents, even though the age and perhaps even the existence of the oldest suggested supercontinent remains

a matter of debate (Bleeker 2003). Continental collision requires convergence and a role for plate tectonics (but see Bédard et al. 2013 for an alternative mechanism), and a similar story is structurally available. Rectilinear, thrust-bound belts characterize late Archean terranes, and in the Pilbara, they date from ~3.1 Ga (Smithies et al. 2007, Van Kranendonk et al. 2007). Large-scale thrust faults, reflecting horizontal compression and dated approximately 2.7 Ga, have been mapped in West Greenland (Friend & Nutman 2005), and a seismic reflection image of an inferred paleosubduction zone has been ascribed to the 2.7 Ga docking of the northern Abitibi granite-greenstone and Opatica plutonic belts, Superior craton Canada (Calvert 1995), both of which are consistent with the commencement of plate tectonics by this time (Cawood et al. 2006).

Geochemical studies offer insights into the conditions of melt generation and the tectonic setting in which new crust was generated. Mafic rocks show a marked increase in La/Yb at \sim 2.7 Ga, a secular decrease in compatible element contents, and an increase in incompatible element contents that reflect a decrease in the degrees of mantle melting through time (Keller & Schoene 2012). Archean terrains are marked by a strongly bimodal distribution in silica (Hawkesworth et al. 2016, Kamber 2015), and this is less well developed in younger areas. Archean felsic rocks are characterized by higher Na₂O/K₂O and La/Yb and less marked Eu anomalies than younger rocks (see Taylor & McLennan 1985). These differences are attributed to partial melting of mafic source rocks at depths in the stability field of garnet rather than plagioclase, likely in response to over thickening of the mafic crust (Kamber 2015, Kemp et al. 2010). Collerson & Kamber (1999) noted an increase in Nb/U and Nb/Th in depleted mantle-derived rocks at ~3 Ga, which they linked with the onset of subduction at that time and the extraction of low Nb/U and Nb/Th material.

An alternative approach is to reconstruct the composition of juvenile continental crust using the initial isotope ratios of preserved continental rocks of different ages. Dhuime et al. (2015) note a marked increase in the Rb/Sr ratio of inferred new crust, and hence of its silica content, at about 3 Ga (**Figure 7**). Prior to 3 Ga, the Rb/Sr ratios of new continental crust are highly scattered, but the median is ~0.03, and it increases to a maximum value of ~0.08 from 3.0 to 1.7 Ga before decreasing to values of ~0.065 in approximately the last 1 Ga. There is a strong positive correlation between Rb/Sr and the SiO₂ contents of crustal rocks and in recent geological settings with the thickness of the continental crust (**Figure 7**) (Dhuime et al. 2015). It was concluded that the SiO₂ contents of new continental crust increased from ~48% before 3.0 Ga to more intermediate compositions in younger terrains with SiO₂ up to ~57%.

The Rb/Sr ratios, and SiO₂ contents, of recent magmatic rocks increase with crustal thickness in Central and South America, and the increase in Rb/Sr appears to be linked to increased differentiation in areas of thicker continental crust (Dhuime et al. 2015). Although the rate of increase of Rb/Sr with crustal thickness may have been different earlier in Earth history, the progressive increase in the time-integrated Rb/Sr ratios of the new crust from 3 Ga to 1 Ga (Figure 7) is taken to reflect at least the relative thickening of the continental crust through time from ~ 20 km at 3 Ga to \sim 40 km by 1 Ga, before it decreases to \sim 30 km toward the present. Assuming that thickness is a proxy for crustal volume, the decrease in inferred crustal volumes presumably reflects crust destruction rates being higher than crust generation rates, perhaps since the onset of cold subduction (Figure 3a) (Brown 2007, 2014). The inferred increase in crustal thickness from 3 Ga to 1 Ga is accompanied by a global increase in the sedimentary contribution to the magmatic record reflected in higher δ^{18} O since the late Archean (Valley et al. 2005) and by increasing 87 Sr/ 86 Sr isotope ratios in seawater from 3 to 2 Ga (Shields 2007). It therefore indicates when significant volumes of continental crust emerged (Arndt 1999; Flament et al. 2008, 2013; Pons et al. 2013; Rey & Coltice 2008) and were available for surficial weathering with the associated drawdown of CO₂ and modification of the Earth's atmosphere (Kramers 2002, Lee et al. 2016).



The estimated Rb/Sr ratios of juvenile continental crust plotted against the age of crust formation for \sim 13,000 whole-rock analyses (Dhuime et al. 2015). Rb/Sr increases with both whole-rock silica content and crustal thickness at the site of magma generation. The calculated volumes of continental crust from Dhuime et al. (2012) are presented for comparison. Stages of Earth evolution: Stage 1 is the early accretion of Earth, which may have persisted for 5–10 Myr (Elkins-Tanton 2008); stage 2 is dominated by vertical tectonics; stage 3 is dominated by hot subduction, stage 4 is Earth's middle age; and stage 5 is characterized by cold subduction.

It is noteworthy that magmatism with within-plate and subduction-like chemical signatures occurred at different times in different Archean areas, and a temporal shift from within-plate to subduction-related magmatism is better developed in some areas than others. However, the changes in the rates of growth of continental crust (**Figure 6**), the ages of the oldest mantle eclogitic samples (**Figure 2**), and the increasing emergence of the continental crust (**Figures 3** and 7) are all taken to reflect the dominant role of plate tectonics by \sim 3 Ga. Subduction-related magmatism occurred before 3 Ga, and new continental crust has been generated in within-plate settings since 3 Ga, such that the bulk composition of the continental crust may contain a \sim 20% contribution of material generated in intraplate settings (see discussion in Rudnick 1995). The first preserved record of a subaerial, large igneous province-related magmatism occurs approximately 3 Ga and becomes widespread by 2.5 Ga (Kump & Barley 2007).

6.3. Preliminary Model for a Two-Phase Evolution of the Continental Crust

Key aspects of the model for the evolution of the continental crust outlined in this contribution are that (*a*) the composition of new continental crust (Dhuime et al. 2015) and the upper crust (Tang et al. 2016) was more mafic before 3 Ga than subsequently; (*b*) at \sim 3 Ga the volume of continental crust was \sim 70% of its present volume; (*c*) there was a significant change in the growth rate of the continental crust at \sim 3 Ga, which has been attributed to the onset of plate tectonics



Schematic illustration of a preliminary two-part model for the generation and evolution of the continental crust. Crust A is the (more mafic) crust generated before 3 Ga (**Figure 7**), and crust B is the more differentiated crust generated subsequently. The simple box model starts with the continental crust growth curve (inset, *blue curve*; Dhuime et al. 2012), and crust destruction rates for crusts A and B were calculated for 500-Myr increments to match the present-day age distribution of juvenile crust (*red curve*; Condie & Aster 2010). The crust generation rates are held constant, and the calculated changes in crust destruction rates for crust A and B are shown.

as the dominant tectonic regime; and (d) less than 5% of the present volume of the continental crust consists of rocks older than 3 Ga.

These observations are incorporated in a simple two-part model of evolution for the continental crust (Figure 8) (Dhuime et al. 2016). This model is not unique, but it illustrates some of the implications that need to be considered. The continental crust generation rates were taken to be constant (e.g., Dhuime et al. 2012), but the composition of the crust changed with time. Juvenile continental crust, prior to 3 Ga, was dominantly mafic (crust A), and from \sim 3 Ga, juvenile crust became more differentiated in composition (crust B), as shown by the inferred increase in the SiO_2 content of juvenile crust plotted in Figure 7. This change is linked to the global onset of plate tectonics as the dominant setting of crust generation and the transition to a modern style of more differentiated continental crust associated with convergent margin settings. In the current model, the crust destruction rates before 3 Ga are taken to be $\sim 10\%$ of those at the present day, and these increased once plate tectonics became the dominant setting of crust generation at 3 Ga. The model has a high-destruction rate of crust A between 3 and 2.5 Ga to account for the low volume of pre-3 Ga crust preserved now, which is consistent with the predominantly mafic and hence relatively dense composition of the pre-3 Ga crust. This is followed by a rapid decrease in the destruction rates of that early crust and an increase in the destruction rates of crust B progressively from the Archean-Proterozoic transition to reach a maximum value at the present day (Figure 8). The volume of crust destroyed to account for the crustal volume at 3.0 Ga being 70% of the present-day value, and <5% of crust older than 3.0 Ga preserved at the present day, indicates that about twice the present volume of the continental crust has been recycled back into the mantle since the onset of plate tectonics (Hawkesworth et al. 2016).

6.4. Earth's Continental Lithosphere Through Time

The geological record preserved today is fragmentary, biased by tectonic and sedimentary processes, and many segments from old oceanic to early Archean lithosphere are barely represented. There are peaks of mantle depletion (T_{RD}) and zircon U-Pb crystallization ages at ~2.8 Ga and younger (**Figures 3** and **5**). Older ages are recorded in diamonds and in crustal rocks and zircons (**Figure 2**), but there is simply not the record for a meaningful comparison of crustal and mantle processes before ~3 Ga. It is argued that plate tectonics was the dominant tectonic process involved in the generation of the continental crust by ~3 Ga as evidenced in the reduction of crustal growth rate, which is attributed to the increased destruction of continental crust along destructive plate margins; the ages of the oldest eclogitic fragments; and changes from more mafic to more felsic compositions of new continental crust. The supercontinent cycle involving dispersal and assembly of continental crust through the opening and closing of ocean basins, at least as currently understood for Pangea and Gondwana, is driven by plate tectonics. Thus, if the early continental record preserves rocks that predate the start of plate tectonics, it is inferred that these rocks should not display the preservation bias associated with the development of supercontinents.

Based on changes in the character of the preserved rock record through time along with constraints provided by numerical modeling and the secular evolution of the mantle's thermal history, we envisage five main stages in the evolution of Earth. The boundary between each stage corresponds with a change in the character of the lithosphere and the tectonic processes controlling its development (Hawkesworth et al. 2016). These five stages are (*a*) initial accretion of Earth; (*b*) generation of crust prior to 3.0 Ga in a preplate tectonic regime; (*c*) early plate tectonics involving hot subduction at relatively high geothermal gradients and extending until approximately 1.7 Ga; (*d*) Earth's middle age from 1.7 to 0.75 Ga, characterized by apparent lithospheric, environmental, and evolutionary stability; and (*e*) initiation of modern cold subduction (i.e., at relatively low geothermal gradients) at ~0.75 Ga. The change from stage 2 to 3 is also associated with the change in the composition of new continental crust (**Figure 7**).

Stage 1 included the initial accretion of Earth, the differentiation of the core and mantle, the development of a magma ocean, and development of an undifferentiated mafic protocrust. It may have persisted for 5-10 Myr (Elkins-Tanton 2008). The second stage extended until approximately 3 Ga. Little lithospheric record remains of the first 600 Myr owing to widespread meteorite bombardment, including collision between the proto-Earth and a body inferred to be approximately the size of Mars and sometimes referred to as Theia, which resulted in the formation of the Moon (Jacobson et al. 2014, Young et al. 2016). The significant scale and frequency of meteorite impact terminated with what is referred to as the end of the late heavy bombardment at approximately 3.9 Ga (Gomes et al. 2005, Marchi et al. 2014). Potential mantle temperatures are inferred to have been \sim 250°C hotter than the present-day average convecting asthenospheric temperature of $\sim 1,350^{\circ}$ C at the end of the late heavy bombardment of meteorites (Herzberg et al. 2010, Korenaga 2013). Numerical modeling suggests widespread asthenospheric melting and impregnation of the lithosphere at such temperatures, reducing lithospheric rigidity and inhibiting, and perhaps preventing, development of sustained deep-level lithospheric subduction (Sizova et al. 2010, van Hunen & van den Berg 2008). Zones of mantle upwelling and melting constituted sites of lithospheric generation, with recycling back into the mantle inferred to occur

through lithospheric delamination (Fischer & Gerya 2016, Johnson et al. 2014, Sizova et al. 2015).

The lack of a garnet signature in the heavy REE variations in residual peridotites and the profiles of olivine composition versus depth for different cratonic lithosphere sections in which the Mg number tends to decrease with increasing depth are taken to reflect melting of a relatively hot mantle at shallow levels (e.g., Canil 2004, Carlson et al. 2005, Helmstaedt & Schulze 1989, Herzberg & Rudnick 2012, Kelemen et al. 1998, Lee et al. 2011, Simon et al. 2007). This combination of high temperatures and shallow depths suggests that melting took place beneath the oceans, that the complementary oceanic crust was destroyed, and that convergent margin processes were involved in the accretion of the residual mantle material to the continental lithosphere, presumably at the time of, or soon after, the stabilization of the overlying crust. It further suggests that there may be no simple relationship between the mantle lithosphere and the overlying crust, at least at this time (see also Sizova et al. 2015).

It is striking that the degrees of melting invoked to generate the residual mantle lithosphere (25–40%) (Herzberg & Rudnick 2012, Walter 1998) are significantly greater than those required to generate the fractionated Lu/Hf and Sm/Nd ratios of mafic continental crust (as subsequently sampled in the magmas from which zircons crystallized), which were $\sim 15\%$ for a simple batch melting model. We infer that the higher degrees of melting took place under relatively thin lithospheric lids in the oceans, whereas the lower degrees of melting involved in the generation of juvenile mafic continental crust reflect melt generation beneath thicker lids in continental areas. The lack of significant Rb/Sr fractionation of Lu/Hf and Sm/Nd, appears to have continued until 3 Ga, whereupon the composition of new crust became more differentiated (**Figure 7**), and the degrees of melting involved in the generation of new continental crust decreased.

The presence of surface water on Earth before 4 Ga has been inferred from slightly elevated δ^{18} O values in Hadean detrital zircon (Cavosie et al. 2005, Harrison 2009, Valley et al. 2002, Wilde et al. 2001). Water-lain sediments in the Isua greenstone belt (Greenland) attest to the presence of oceans by at least 3.7 Ga (Bolhar et al. 2004). Marine carbonates deposited at 3.48 Ga in the Pilbara craton (Western Australia) preserve evidence for cyanobacteria (Van Kranendonk et al. 2008), which implies photosynthesis and hence start of oxygenation of the atmosphere.

The stabilization of Archean cratons and the change in crustal growth rates at about 3.0 Ga are taken to mark the onset of stage 3, which extended until approximately 1.7 Ga. Mantle temperatures had cooled sufficiently to enable subduction of coherent semirigid plates (Sizova et al. 2010); water was returned to the mantle, which may in turn have facilitated plate tectonics (Korenaga 2013); and the composition of juvenile crust had become more intermediate in composition (Figure 7). Plate tectonics became the dominant regime in which new continental crust was generated (Figure 9), the continental crust became thicker and more evolved, supercontinent cycles developed, and there was increased erosion to the oceans due to thickening and subaerial exposure of continental crust resulting in an increase in Sr isotope ratios in seawater (Figure 3). If \sim 70% of the present volume of the continental crust was present at 3.0 Ga, and <5% are preserved at the present day (Figure 6), it follows that large volumes of continental crust have been destroyed, and yet the overall net growth of crust requires the rate of crust generation to exceed that of recycling (Figure 8). The change from a nonplate tectonic Earth to one in which plate tectonics dominated was unlikely to have been abrupt (Section 6.2). If (a) the inferred changes in composition and thickness of the lithosphere and (b) the linked changes in the surficial Earth such as ocean composition are correct and correspond to secular mantle cooling, then the transition from a predominantly nonplate tectonic Earth is likely to have extended over hundreds of millions of years about 3.0 Ga. Furthermore, rock relations



Schematic depiction of changing tectonic processes controlling the evolution of the lithosphere from an early Earth dominated by nonplate processes (stage 2) to one in which plate tectonics is the main mechanism for the generation and recycling of lithosphere (stages 3–5). These changes are a response to the secular cooling of the mantle and the consequent increase in lithospheric strength and rigidity. The mantle xenolith record suggests that they represent residues (from relatively shallow partial melting of hot mantle beneath the oceans), which subsequently accreted beneath continents. In contrast, the mafic crust that is the source of Archean tonalite-trondhjemite-granodiorites has relatively low Lu/Hf and Sm/Nd ratios, implying lower degrees of partial melting.

similar to those in modern convergent plate margins in some regions suggest that at least localized subduction commenced even earlier (Jenner et al. 2009, O'Neil et al. 2011, Turner et al. 2014).

The onset of plate tectonics resulted in significant changes in the nature of Earth processes preserved in the rock archive during the remainder of stage 3. Oxygen levels in the atmosphere increased significantly in the early Proterozoic during the Great Oxidation Event (Holland 2006, Kump 2008 and references therein). A rise in the atmospheric oxygen level followed on from significant increases in continental lithosphere volume and thickness, the change to a more felsic continental crustal composition, and widespread exposure of the continental crust. This is consistent with a causative relationship, and recently Lee et al. (2016) have linked these changes to a decrease in the oxidative efficiency of Earth's surface, allowing a rise in O₂ levels. Global compilations of δ^{18} O values (**Figure 3**) in zircon display a pattern of peaks and troughs coincident with zircon U-Pb age distribution patterns, except for the ~2.7 Ga age peak, which has only slightly elevated δ^{18} O values (Dhuime et al. 2012, Spencer et al. 2014). The deviation of δ^{18} O values from mantle values is taken to reflect increased incorporation of high δ^{18} O sediment into igneous zircons during crustal reworking and thickening, associated with periods of continental collision and supercontinent assembly, especially since the beginning of the Proterozoic.

The fourth stage is from 1.7–0.75 Ga, recently referred to as Earth's middle age (Cawood & Hawkesworth 2014) and before that as the boring billion (Holland 2006). It is characterized by an absence of significant anomalies in the paleoseawater Sr isotope record and in Hf isotopes in detrital zircon (**Figure 3**), indicating that it was a period of environmental and lithospheric stability (Brasier & Lindsay 1998), and there is a paucity of preserved passive margins (Bradley 2008). This stability may have been linked to the persistence of a relatively stable continental assemblage initiated during assembly of the Nuna supercontinent by ~1.7 Ga and until breakup of its successor, Rodinia, approximately 0.75 Ga (Cawood & Hawkesworth 2014, Cawood et al.

2016). This in turn marks the onset of stage 5 and the development of cold subduction recognized by the first appearance of high- to ultrahigh-pressure metamorphic rocks (Brown 2006, 2014). Falling mantle temperatures, to no more than 80–100°C greater than today, enabled deeper levels of slab break-off in collision zones and the resultant greater depths to which continental crust was subducted prior to exhumation, allowing the development of ultrahigh-pressure metamorphic assemblages (Brown 2006, Sizova et al. 2014). Stage 5 is marked by a strongly episodic distribution of ages linked to the supercontinent cycles of Gondwana and Pangea (**Figure 3**). The estimated thickness, and by analogy the volume, of continental crust decreased perhaps because crust was more efficiently destroyed after the Neoproterozoic transition to the modern plate tectonics regime at relatively low geothermal gradients (Brown 2007). Oxygen levels in both the atmosphere and deep oceans increased, and phosphate and evaporite deposits became widespread, providing a major stimulus for metazoans evolution.

7. WAYS FORWARD

Major developments in geochemical techniques have resulted in a large number of high-quality in situ analyses for radiogenic and stable isotopes in mineral archives that yield precise crystallization ages. High-quality U-Pb zircon analyses, together with T_{RD} Os isotope ages, have allowed for improved statistical approaches in the analysis of variations in the age and composition of the continental crust and its mantle lithosphere and provided insight into the secular evolution of processes involved in the generation and recycling of the lithosphere. There are, however, concerns over biases in the composition and the ages of the rocks preserved, and this highlights the longstanding issue of how best to evaluate the global context of more regional case studies. This remains of particular concern for studies of Archean geology given the relatively small volume of rock that is preserved, but one way to put regional case studies in perspective is in the development of global models of, for example, rates of crustal growth. There is evidence for subduction-related magmatism before 3 Ga, and clearly for within-plate magmatism subsequently, yet the inferred change in global crustal growth rates can be used to argue that plate tectonics became the dominant framework for the generation of continental crust from ~3 Ga. Geodynamic models may also have a particular role as they allow us to make general predictions that are difficult to derive from the very limited (and potentially unrepresentative) rock record.

Other aspects to highlight include improved methods to evaluate the volume of lessdifferentiated continental crust, i.e., that contains little or no zircon, and the composition and the tectonic setting of new continental crust, for example, by evaluation of the Rb/Sr and the U/Pb ratios in the sources of crustal melts (Delavault et al. 2016, Dhuime et al. 2015). Given the concerns of how representative the rock record may be, and the biases present in the sedimentary record, there is a clear need for better constraints on a global value for the erosion parameter K (Allègre & Rousseau 1984) and for improved ways to sample well-mixed reservoirs, such as the upper mantle and the oceans. Improved resolution of mantle anisotropy offers new insights into possible structural differences between different cratonic areas and perhaps the timing of the development of such fabric in the upper mantle (e.g., Adam & Lebedev 2012, Kaczmarek et al. 2016). These approaches may determine the extent to which there are tectonic links between the continental mantle lithosphere and the overlying crust (Griffin et al. 2009, Pearson & Wittig 2013). There is increasing agreement that at least some of the rock record reflects conditions before plate tectonics was the dominant tectonic setting in which new continental crust was generated and reflects the nature of that transition. This too appears to be a fertile area to integrate geodynamic modeling and well-constrained data from the rock record.

DISCLOSURE STATEMENT

The authors are not aware of any affiliations, memberships, funding, or financial holdings that might be perceived as affecting the objectivity of this review.

ACKNOWLEDGMENTS

We thank John Dewey, Mike Daly, Mike Kendall, Tony Prave, Ruth Robinson, Mike Walter, and Tony Watts for ongoing discussions regarding the evolution of Earth's lithosphere, and thank Graham Pearson for providing an updated file on Os isotope data on mantle kimberlites from cratonic areas. Ian Fitzsimons, Carol Frost, Janet Hergt, Leigh Royden, and Simon Turner kindly provided comments on an earlier draft. We acknowledge funding support from the Natural Environment Research Council (grants NE/J021822/1 and NE/K008862/1), the Leverhulme Trust (grant RPG-2015–422), and the Australian Research Council (FL160100168 to P.A.C. and FT100100059 to T.I.S.K.).

LITERATURE CITED

Adam JM-C, Lebedev S. 2012. Azimuthal anisotropy beneath southern Africa from very broad-band surfacewave dispersion measurements. *Geophys. J. Int.* 191(1):155–74

Albarède F. 1998. The growth of continental crust. Tectonophysics 296(1-2):1-14

- Allan C. 2014. Is there a bias between landscape relief and model age? BSc Thesis (Hons), Univ. St Andrews
- Allègre CJ, Hart SR, Minster JF. 1983. Chemical structure and evolution of the mantle and continents determined by inversion of Nd and Sr isotopic data, I. Theoretical methods. *Earth Planet. Sci. Lett.* 66(0):177–90
- Allègre CJ, Rousseau D. 1984. The growth of the continent through geological time studied by Nd isotope analysis of shales. *Earth Planet. Sci. Lett.* 67(1):19–34
- Anhaeusser CR, Robb LJ. 1981. Magmatic cycles and the evolution of the Archaean granitic crust in the eastern Transvaal and Swaziland. Spec. Publ., Geol. Soc. Aust. 7:457–67
- Armstrong RL. 1981. Radiogenic isotopes: the case for crustal recycling on a steady-state no-continentalgrowth Earth. Philos. Trans. R. Soc. Lond. Ser. A 301:443–72
- Arndt N. 1999. Why was flood volcanism on submerged continental platforms so common in the Precambrian? Precambrian Res. 97(3–4):155–64
- Arndt N, Davaille A. 2013. Episodic Earth evolution. Tectonophysics 609:661-74
- Arndt NT, Goldstein SL. 1987. Use and abuse of crust-formation ages. Geology 15(10):893-95
- Artemieva IM. 2009. The continental lithosphere: reconciling thermal, seismic, and petrologic data. *Lithos* 109(1-2):23-46
- Artemieva IM, Mooney WD. 2001. Thermal thickness and evolution of Precambrian lithosphere: a global study. J. Geophys. Res. 106(B8):16387–414
- Becker H, Horan MF, Walker RJ, Gao S, Lorand JP, Rudnick RL. 2006. Highly siderophile element composition of the Earth's primitive upper mantle: constraints from new data on peridotite massifs and xenoliths. *Geochim. Cosmochim. Acta* 70(17):4528–50
- Becker H, Shirey SB, Carlson RW. 2001. Effects of melt percolation on the Re-Os systematics of peridotites from a Paleozoic convergent plate margin. *Earth Planet. Sci. Lett.* 188(1–2):107–21
- Bédard JH, Harris LB, Thurston PC. 2013. The hunting of the snArc. Precambrian Res. 229:20-48
- Belousova EA, Kostitsyn YA, Griffin WL, Begg GC, O'Reilly SY, Pearson NJ. 2010. The growth of the continental crust: constraints from zircon Hf-isotope data. *Lithos* 119(3–4):457–66
- Bindeman IN, Bekker A, Zakharov DO. 2016. Oxygen isotope perspective on crustal evolution on early Earth: a record of Precambrian shales with emphasis on Paleoproterozoic glaciations and Great Oxygenation Event. *Earth Planet. Sci. Lett.* 437:101–13
- Bleeker W. 2003. The late Archean record: a puzzle in ca. 35 pieces. Lithos 71(2-4):99-134

- Bodinier JL, Godard M. 2003. Orogenic, ophiolitic, and abyssal peridotites. In *Treatise on Geochemistry*, Vol. 2: *The Mantle and Core*, ed. KK Turekian, HD Holland, pp. 1–73. New York: Elsevier
- Bolhar R, Kamber BS, Moorbath S, Fedo CM, Whitehouse MJ. 2004. Characterisation of early Archaean chemical sediments by trace element signatures. *Earth Planet. Sci. Lett.* 222(1):43–60
- Bowring AS, Williams SI. 1999. Priscoan (4.00–4.03 Ga) orthogneisses from northwestern Canada. Contrib. Mineral. Petrol. 134(1):3–16
- Boyd FR. 1989. Compositional distinction between oceanic and cratonic lithosphere. *Earth Planet. Sci. Lett.* 96(1–2):15–26
- Boyd FR, Pokhilenko NP, Pearson DG, Mertzman SA, Sobolev NV, Finger LW. 1997. Composition of the Siberian cratonic mantle: evidence from Udachnaya peridotite xenoliths. *Contrib. Mineral. Petrol.* 128:228–46
- Bradley DC. 2008. Passive margins through earth history. Earth-Sci. Rev. 91(1-4):1-26
- Bradley DC. 2011. Secular trends in the geologic record and the supercontinent cycle. *Earth-Sci. Rev.* 108(1–2):16–33
- Brasier MD, Lindsay JF. 1998. A billion years of environmental stability and the emergence of eukaryotes: new data from northern Australia. *Geology* 26(6):555–58
- Brown M. 2006. Duality of thermal regimes is the distinctive characteristic of plate tectonics since the Neoarchean. *Geology* 34(11):961–64
- Brown M. 2007. Metamorphic conditions in orogenic belts: a record of secular change. *Int. Geol. Rev.* 49(3):193–234
- Brown M. 2014. The contribution of metamorphic petrology to understanding lithosphere evolution and geodynamics. *Geosci. Front.* 5(4):553–69
- Burton KW, Schiano P, Birck JL, Allègre CJ, Rehkämper M, et al. 2000. The distribution and behaviour of rhenium and osmium amongst mantle minerals and the age of the lithospheric mantle beneath Tanzania. *Earth Planet. Sci. Lett.* 183(1–2):93–106
- Calvert AJ. 1995. Seismic evidence for a magma chamber beneath the slow-spreading Mid-Atlantic Ridge. *Nature* 377:410–14
- Campbell IH. 2003. Constraints on continental growth models from Nb/U ratios in the 3.5 Ga Barberton and other Archaean basalt-komatiite suites. *Am. J. Sci.* 303(4):319–51
- Campbell IH, Allen CM. 2008. Formation of supercontinents linked to increases in atmospheric oxygen. *Nat. Geosci.* 1:554–58
- Campbell IH, Squire RJ. 2010. The mountains that triggered the Late Neoproterozoic increase in oxygen: the Second Great Oxidation Event. *Geochim. Cosmochim. Acta* 74(15):4187–206
- Canil D. 2004. Mildly incompatible elements in peridotites and the origins of mantle lithosphere. *Lithos* 77(1-4):375-93
- Carlson RW, Irving AJ. 1994. Depletion and enrichment history of subcontinental lithospheric mantle: an Os, Sr, Nd and Pb isotopic study of ultramafic xenoliths from the northwestern Wyoming Craton. *Earth Planet. Sci. Lett.* 126(4):457–72
- Carlson RW, Irving AJ, Schulze DJ, Hearn BC Jr. 2004. Timing of Precambrian melt depletion and Phanerozoic refertilization events in the lithospheric mantle of the Wyoming Craton and adjacent Central Plains Orogen. *Lithos* 77(1–4):453–72
- Carlson RW, Pearson DG, James DE. 2005. Physical, chemical, and chronological characteristics of continental mantle. *Rev. Geophys.* 43(1):RG1001
- Cavosie AJ, Valley JW, Wilde SA. 2005. Magmatic 8¹⁸O in 4400–3900 Ma detrital zircons: a record of the alteration and recycling of crust in the Early Archean. *Earth Planet. Sci. Lett.* 235(3–4):663–81
- Cawood PA, Buchan C. 2007. Linking accretionary orogenesis with supercontinent assembly. *Earth-Sci. Rev.* 82:217–56
- Cawood PA, Hawkesworth CJ. 2014. Earth's middle age. Geology 42(6):503-6
- Cawood PA, Hawkesworth CJ, Dhuime B. 2013. The continental record and the generation of continental crust. *Geol. Soc. Am. Bull.* 125(1–2):14–32
- Cawood PA, Kröner A, Pisarevsky S. 2006. Precambrian plate tectonics: criteria and evidence. GSA Today 16:4–11

- Cawood PA, Nemchin AA, Freeman MJ, Sircombe KN. 2003. Linking source and sedimentary basin: detrital zircon record of sediment flux along a modern river system and implications for provenance studies. *Earth Planet. Sci. Lett.* 210:259–68
- Cawood PA, Strachan RA, Pisarevsky SA, Gladkochub DP, Murphy JB. 2016. Linking collisional and accretionary orogens during Rodinia assembly and breakup: implications for models of supercontinent cycles. *Earth Planet. Sci. Lett.* 449:118–26
- Chauvel C, Garçon M, Bureau S, Besnault A, Jahn B-M, Ding Z. 2014. Constraints from loess on the Hf-Nd isotopic composition of the upper continental crust. *Earth Planet. Sci. Lett.* 388:48–58
- Chesley JT, Rudnick RL, Lee C-T. 1999. Re-Os systematics of mantle xenoliths from the East African Rift: age, structure, and history of the Tanzanian craton. *Geochim. Cosmochim. Acta* 63(7–8):1203–17
- Choi SH, Suzuki K, Mukasa SB, Lee J-I, Jung H. 2010. Lu-Hf and Re-Os systematics of peridotite xenoliths from Spitsbergen, western Svalbard: implications for mantle-crust coupling. *Earth Planet. Sci. Lett.* 297(1– 2):121–32
- Clift PD, Vannucchi P, Morgan JP. 2009. Crustal redistribution, crust-mantle recycling and Phanerozoic evolution of the continental crust. *Earth-Sci. Rev.* 97(1–4):80–104
- Cloos M. 1993. Lithospheric buoyancy and collisional orogenesis: subduction of oceanic plateaus, continental margins, island arcs, spreading ridges, and seamounts. *Geol. Soc. Am. Bull.* 105(6):715–37
- Collerson KD, Kamber BS. 1999. Evolution of the continents and the atmosphere inferred from Th-U-Nb systematics of the depleted mantle. *Science* 283(5407):1519–22
- Condie KC. 1998. Episodic continental growth and supercontinents: a mantle avalanche connection? *Earth Planet. Sci. Lett.* 163(1–4):97–108
- Condie KC, Aster RC. 2010. Episodic zircon age spectra of orogenic granitoids: the supercontinent connection and continental growth. *Precambrian Res.* 180(3–4):227–36
- Condie KC, Davaille A, Aster RC, Arndt N. 2015. Upstairs-downstairs: Supercontinents and large igneous provinces, are they related? *Int. Geol. Rev.* 57(11–12):1341–48
- Dana JD. 1873. On some results of the earth's contraction from cooling, including a discussion of the origin of mountains, and the nature of the earth's interior. *Am. J. Sci., Ser.* 3. 5:423–43
- Davidson JP, Arculus RJ. 2006. The significance of Phanerozoic arc magmatism in generating continental crust. In *Evolution and Differentiation of the Continental Crust*, ed. M Brown, T Rushmer, pp. 135–72. Cambridge, UK: Cambridge Univ. Press
- Delavault H, Dhuime B, Hawkesworth CJ, Cawood PA, Marschall H, Edinburgh Ion Microprobe Facility. 2016. Tectonic settings of continental crust formation: Insights from Pb isotopes in feldspar inclusions in zircon. *Geology* 44(10):819–22
- Dewey JF, Windley BF. 1981. Growth and differentiation of the continental crust. Philos. Trans. R. Soc. Lond. Ser. A 301(1461):189–206
- Dhuime B, Hawkesworth CJ, Cawood PA, Storey CD. 2012. A change in the geodynamics of continental growth 3 billion years ago. *Science* 335(6074):1334–36
- Dhuime B, Hawkesworth CJ, Storey CD, Cawood PA. 2011. From sediments to their source rocks: Hf and Nd isotopes in recent river sediments. *Geology* 39(4):407–10
- Dhuime B, Hawkesworth CJ, Wookey J. 2016. Secular evolution of the rates of generation and destruction of the continental crust. Presented at Goldschmidt Conf., June 26–July 1, Yokohama, Jpn. (Abstr. 660)
- Dhuime B, Wuestefeld A, Hawkesworth CJ. 2015. Emergence of modern continental crust about 3 billion years ago. Nat. Geosci. 8(7):552–55
- Elkins-Tanton LT. 2008. Linked magma ocean solidification and atmospheric growth for Earth and Mars. *Earth Planet. Sci. Lett.* 271(1–4):181–91
- Evans DAD, Pisarevsky SA. 2008. Plate tectonics on early Earth? Weighing the paleomagnetic evidence. Geol. Soc. Am. Spec. Pap. 440: 249–63
- Farquhar J, Zerkle AL, Bekker A. 2013. Geologic and geochemical constraints on the earth's early atmosphere. In *Treatise on Geochemistry: Reference Module in Earth Systems and Environmental Sciences*, ed. KK Turekian, HD Holland, pp. 91–138. New York: Elsevier. 2nd ed.
- Finnerty AA, Boyd FR. 1987. Thermobarometry for garnet peridotites: basis for the determination of thermal and compositional structure of the upper mantle. In *Mantle Xenoliths*, ed. PH Nixon, pp. 381–402. Hoboken, NJ: Wiley

- Fischer R, Gerya T. 2016. Regimes of subduction and lithospheric dynamics in the Precambrian: 3D thermomechanical modelling. *Gondwana Res.* 37:53–70
- Flament N, Coltice N, Rey PF. 2008. A case for late-Archaean continental emergence from thermal evolution models and hypsometry. *Earth Planet. Sci. Lett.* 275(3–4):326–36
- Flament N, Coltice N, Rey PF. 2013. The evolution of the ⁸⁷Sr/⁸⁶Sr of marine carbonates does not constrain continental growth. *Precambrian Res.* 229(0):177–88
- Friend CRL, Nutman AP. 2005. New pieces to the Archaean terrane jigsaw puzzle in the Nuuk region, southern West Greenland: steps in transforming a simple insight into a complex regional tectonothermal model. J. Geol. Soc. Lond. 162(1):147–62
- Gao S, Rudnick RL, Carlson RW, McDonough WF, Liu Y-S. 2002. Re-Os evidence for replacement of ancient mantle lithosphere beneath the North China craton. *Earth Planet. Sci. Lett.* 198(3–4):307–22

Garrels RM, Mackenzie FT. 1971. Evolution of Sedimentary Rocks. New York: Norton

- Gastil RG. 1960. The distribution of mineral dates in time and space. Am. J. Sci. 258(1):1-35
- Goldfarb RJ, Groves DI, Gardoll S. 2001. Orogenic gold and geologic time: a global synthesis. Ore Geol. Rev.18:1–75
- Goldstein SJ, Jacobsen SB. 1988. Nd and Sr isotopic systematics of river water suspended material: implications for crustal evolution. *Earth Planet. Sci. Lett.* 87(3):249–65
- Gomes R, Levison HF, Tsiganis K, Morbidelli A. 2005. Origin of the cataclysmic Late Heavy Bombardment period of the terrestrial planets. *Nature* 435(7041):466–69
- Goodwin AM. 1996. Principles of Precambrian Geology. London: Academic
- Gower CF, Krogh TE. 2002. A U-Pb geochronological review of the Proterozoic history of the eastern Grenville Province. Can. J. Earth Sci. 39:795–829
- Grand SP. 2002. Mantle shear-wave tomography and the fate of subducted slabs. *Philos. Trans. R. Soc. A* 360(1800):2475–91
- Griffin WL, O'Reilly SY, Afonso JC, Begg GC. 2009. The composition and evolution of lithospheric mantle: a re-evaluation and its tectonic implications. *J. Petrol.* 50(7):1185–204
- Hacker BR, Kelemen PB, Behn MD. 2015. Continental lower crust. Annu. Rev. Earth Planet. Sci. 43:167-205
- Handler MR, Bennett VC, Carlson RW. 2005. Nd, Sr and Os isotope systematics in young, fertile spinel peridotite xenoliths from northern Queensland, Australia: a unique view of depleted MORB mantle? *Geochim. Cosmochim. Acta* 69(24):5747–63
- Handler MR, Bennett VC, Esat TM. 1997. The persistence of off-cratonic lithospheric mantle: Os isotopic systematics of variably metasomatised southeast Australian xenoliths. *Earth Planet. Sci. Lett.* 151(1–2):61– 75
- Handler MR, Wysoczanski RJ, Gamble JA. 2003. Proterozoic lithosphere in Marie Byrd Land, West Antarctica: Re-Os systematics of spinel peridotite xenoliths. *Chem. Geol.* 196(1–4):131–45
- Hansen VL. 2015. Impact origin of Archean cratons. Lithosphere 7(5):563-78
- Harrison TM. 2009. The Hadean crust: evidence from >4 Ga zircons. Annu. Rev. Earth Planet. Sci. 37:479-505
- Hawkesworth C, Cawood P, Kemp T, Storey C, Dhuime B. 2009. A matter of preservation. Science 323:49-50
- Hawkesworth CJ, Cawood PA, Dhuime B. 2016. Tectonics and crustal evolution. GSA Today 26(9):4-11
- Helmstaedt H, Schulze DJ. 1989. Southern African kimberlites and their mantle sample: implications for Archean tectonics and lithospheric evolution. In *Kimberlites and Related Rocks*, Vol. 1: *Their Composition*, *Occurrence*, *Origin and Emplacement*, ed. J Ross, pp. 358–68. Sydney: Geol. Soc. Aust.
- Herzberg C, Condie K, Korenaga J. 2010. Thermal history of the Earth and its petrological expression. Earth Planet. Sci. Lett. 292(1–2):79–88
- Herzberg C, Rudnick R. 2012. Formation of cratonic lithosphere: an integrated thermal and petrological model. *Lithus* 149(0):4–15
- Holland HD. 2006. The oxygenation of the atmosphere and oceans. Philos. Trans. R. Soc. Lond. Ser. B 361(1470):903-15
- Hurley PM, Hughes H, Faure G, Fairbairn HW, Pinson WH. 1962. Radiogenic strontium-87 model of continent formation. 7. Geophys. Res. 67(13):5315-34
- Hutton J. 1785. Abstract of a dissertation read in the Royal Society of Edinburgh, upon the seventh of March, and fourth of April, MDCCLXXXV, concerning the System of the Earth, its duration, and stability, R. Soc. Edinb.

Hutton J. 1788. Theory of the Earth; or an investigation of the laws observable in the composition, dissolution, and restoration of land upon the globe. *Trans. R. Soc. Edinb.* 1:209–304

Jacobsen SB. 1988. Isotopic constraints on crustal growth and recycling Earth Planet. Sci. Lett. 90(3):315-29

- Jacobsen SB, Wasserburg GJ. 1979. Nd and Sr isotopic study of the Bay of Islands ophiolite complex and the evolution of the source of midocean ridge basalts. *J. Geophys. Res.* 84(B13):7429–45
- Jacobson SA, Morbidelli A, Raymond SN, O'Brien DP, Walsh KJ, Rubie DC. 2014. Highly siderophile elements in Earth's mantle as a clock for the Moon-forming impact. *Nature* 508(7494):84–87
- James DE, Fouch MJ, VanDecar JC, Lee SVD. 2001. Tectospheric structure beneath southern Africa. Geophys. Res. Lett. 28(13):2485–88
- Janney PE, Shirey SB, Carlson RW, Pearson DG, Bell DR, et al. 2010. Age, composition and thermal characteristics of South African off-craton mantle lithosphere: evidence for a multi-stage history. J. Petrol. 51(9):1849–90
- Jaupart C, Mareschal JC. 2003. Constraints on crustal heat production from heat flow data. In Treatise on Geochemistry, Vol. 3: The Crust, ed. KK Turekian, HD Holland, pp. 65–84. New York: Elsevier
- Jaupart C, Mareschal JC, Guillou-Frottier L, Davaille A. 1998. Heat flow and thickness of the lithosphere in the Canadian Shield. J. Geophys. Res. 103(B7):15269–86
- Jenner FE, Bennett VC, Nutman AP, Friend CRL, Norman MD, Yaxley G. 2009. Evidence for subduction at 3.8 Ga: geochemistry of arc-like metabasalts from the southern edge of the Isua supracrustal belt. *Chem. Geol.* 261(1–2):82–97
- Jenner FE, Bennett VC, Yaxley G, Friend CRL, Nebel O. 2013. Eoarchean within-plate basalts from southwest Greenland. *Geology* 41(3):327–30
- Johnson TE, Brown M, Kaus BJP, VanTongeren JA. 2014. Delamination and recycling of Archaean crust caused by gravitational instabilities. *Nat. Geosci.* 7(1):47–52

Jordan TH. 1978. Composition and development of the continental tectosphere. Nature 274(5671):544-48

- Jordan TH. 1988. Structure and formation of the continental tectosphere. J. Petrol. Spec. Vol. (1):11-37
- Kaczmarek M-A, Reddy SM, Nutman AP, Friend CRL, Bennett VC. 2016. Earth's oldest mantle fabrics indicate Eoarchaean subduction. Nat. Commun. 7:10665
- Kamber BS. 2015. The evolving nature of terrestrial crust from the Hadean, through the Archaean, into the Proterozoic. *Precambrian Res.* 258(0):48–82
- Kearey P, Klepeis KA, Vine FJ. 2009. Global Tectonics. Oxford, UK: Wiley
- Kelemen PB, Hart SR, Bernstein S. 1998. Silica enrichment in the continental upper mantle via melt/rock reaction. *Earth Planet. Sci. Lett.* 164(1–2):387–406
- Keller CB, Schoene B. 2012. Statistical geochemistry reveals disruption in secular lithospheric evolution about 2.5 Gyr ago. Nature 485(7399):490–93
- Kemp AIS, Wilde SA, Hawkesworth CJ, Coath CD, Nemchin A, et al. 2010, Hadean crustal evolution revisited: new constraints from Pb-Hf isotope systematics of the Jack Hills zircons. *Earth Planet. Sci. Lett.* 296(1–2):45–563
- Kober L. 1921. Der Bau der Erde. Berlin: Gebrüder Borntraeger
- Korenaga J. 2013. Initiation and evolution of plate tectonics on Earth: theories and observations. Annu. Rev. Earth Planet. Sci. 41(1):117–51
- Kramers JD. 2002. Global modelling of continent formation and destruction through geological time and implications for CO₂ drawdown in the Archaean Eon. *Geol. Soc. Lond. Spec. Publ.* 199:259–74
- Kramers JD, Tolstikhin IN. 1997. Two terrestrial lead isotope paradoxes, forward transport modelling, core formation and the history of the continental crust. *Chem. Geol.* 139(1–4):75–110
- Kump LR. 2008. The rise of atmospheric oxygen. Nature 451(7176):277-78
- Kump LR, Barley ME. 2007. Increased subaerial volcanism and the rise of atmospheric oxygen 2.5 billion years ago. Nature 448(7157):1033–36
- Larsen IJ, Montgomery DR, Greenberg HM. 2014. The contribution of mountains to global denudation. Geology 42(6):527–30
- Lee C-T, Yin Q, Rudnick RL, Chesley JT, Jacobsen SB. 2000. Osmium isotopic evidence for Mesozoic removal of lithospheric mantle beneath the Sierra Nevada, California. *Science* 289(5486):1912–16
- Lee C-TA, Luffi P, Chin EJ. 2011. Building and destroying continental mantle. *Annu. Rev. Earth Planet. Sci.* 39(1):59–90

- Lee C-TA, Yeung LY, McKenzie NR, Yokoyama Y, Ozaki K, Lenardic A. 2016. Two-step rise of atmospheric oxygen linked to the growth of continents. *Nat. Geosci.* 9(6):417–24
- Lee S, Nolet G. 1997. Upper mantle S velocity structure of North America. J. Geophys. Res. 102(B10):22815–38
- Liu J, Riches AJV, Pearson DG, Luo Y, Kienlen B, et al. 2016, Age and evolution of the deep continental root beneath the central Rae craton, northern Canada. *Precambrian Res.* 272:168–84
- Liu J, Rudnick RL, Walker RJ, Gao S, Wu F-Y, et al. 2011. Mapping lithospheric boundaries using Os isotopes of mantle xenoliths: an example from the North China Craton. *Geochim. Cosmochim. Acta* 75(13):3881– 902
- Liu J, Scott JM, Martin CE, Pearson DG. 2015. The longevity of Archean mantle residues in the convecting upper mantle and their role in young continent formation. *Earth Planet. Sci. Lett.* 424:109–18
- Luguet A, Jaques AL, Pearson DG, Smith CB, Bulanova GP, et al. 2009. An integrated petrological, geochemical and Re-Os isotope study of peridotite xenoliths from the Argyle lamproite, Western Australia and implications for cratonic diamond occurrences. *Lithus* 112(Suppl. 2):1096–108
- Malitch KN, Merkle RKW. 2004. Ru-Os-Ir-Pt and Pt-Fe alloys from the Evander goldfield, Witwatersrand Basin, South Africa: detrital origin inferred from compositional and osmium isotope data. *Can. Mineral.* 42(2):631–50
- Marchi S, Bottke WF, Elkins-Tanton LT, Bierhaus M, Wuennemann K, et al. 2014. Widespread mixing and burial of Earth's Hadean crust by asteroid impacts. *Nature* 511(7511):578–82
- Matsukage KN, Kawasaki T. 2014. Hydrous origin of the continental cratonic mantle. *Earth, Planets Space* 66:29
- McCulloch MT, Wasserburg GJ. 1978. Sm-Nd and Rb-Sr chronology of continental crust formation. *Science* 200:1003–11
- McKenzie D, Daly MC, Priestley K. 2015. The lithospheric structure of Pangea. Geology 43(9):783-86
- McKenzie D, Priestley K. 2008. The influence of lithospheric thickness variations on continental evolution. *Lithos* 102(1–2):1–11
- McLennan SM, Taylor SR, Hemming SR. 2005. Composition, differentiation, and evolution of continental crust: constraints from sedimentary rocks and heat flow. In *Evolution and Differentiation of the Continental Crust*, ed. M Brown, T Rushwer, pp. 93–135. Cambridge, UK: Cambridge Univ. Press
- Meisel T, Walker RJ, Irving AJ, Lorand J-P. 2001. Osmium isotopic compositions of mantle xenoliths: a global perspective. *Geochim. Cosmochim. Acta* 65(8):1311–23
- Mooney WD, Laske G, Masters TG. 1998. CRUST 5.1: A global crustal model at 5° × 5°. *J. Geophys. Res.* 103(B1):727–47
- Moorbath S. 1977. Ages, isotopes and evolution of Precambrian continental crust. Chem. Geol. 20:151-87
- Moores EM, Twiss RJ. 1995. Tectonics. New York: Freeman
- Mosier DL, Berger VI, Singer DA. 2009. Volcanogenic massive sulfide deposits of the world—database and grade and tonnage models. US Geol. Surv. Open-File Rep. 2009-1034. https://pubs.usgs.gov/of/2009/1034/
- Næraa T, Scherstén A, Rosing MT, Kemp AIS, Hoffmann JE, et al. 2012. Hafnium isotope evidence for a transition in the dynamics of continental growth 3.2 Gyr ago. *Nature* 485(7400):627–30
- Nance RD, Murphy JB, Santosh M. 2014. The supercontinent cycle: a retrospective essay. *Gondwana Res.* 25(1):4–29
- Nyblade AA. 1999. Heat flow and the structure of Precambrian lithosphere. Lithos 48(1-4):81-91
- O'Neil J, Francis D, Carlson RW. 2011. Implications of the Nuvvuagittuq greenstone belt for the formation of Earth's early crust. *J. Petrol.* 52(5):985–1009
- O'Nions RK, Evensen NM, Hamilton PJ. 1979. Geochemical modeling of mantle differentiation and crustal growth. J. Geophys. Res. 84(B11):6091–101
- O'Nions RK, Hamilton PJ, Hooker PJ. 1983. A Nd isotope investigation of sediments related to crustal development in the British Isles. *Earth Planet. Sci. Lett.* 63(2):229–40
- Parman SW. 2015. Time-lapse zirconography: imaging punctuated continental evolution. *Geochem. Perspect. Lett.* 1:43–52
- Pearson DG, Carlson RW, Shirey SB, Boyd FR, Nixon PH. 1995. Stabilisation of Archaean lithospheric mantle: a Re-Os isotope study of peridotite xenoliths from the Kaapvaal craton. *Earth Planet. Sci. Lett.* 134(3–4):341–57

- Pearson DG, Wittig N. 2013. The formation and evolution of cratonic mantle lithosphere—evidence from mantle xenoliths. In *Treatise on Geochemistry*, ed. KK Turekian, HD Holland, pp. 255–92. New York: Elsevier. 2nd ed.
- Peslier AH, Reisberg L, Ludden J, Francis D. 2000a. Os isotopic systematics in mantle xenoliths; age constraints on the Canadian Cordillera lithosphere. *Chem. Geol.* 166(1–2):85–101
- Peslier AH, Reisberg L, Ludden J, Francis D. 2000b. Re-Os constraints on harzburgite and lherzolite formation in the lithospheric mantle: a study of northern Canadian Cordillera xenoliths. *Geochim. Cosmochim. Acta* 64(17):3061–71
- Pollack HN. 1986. Cratonization and thermal evolution of the mantle. Earth Planet. Sci. Lett. 80(1-2):175-82
- Pollack HN, Hurter SJ, Johnson JR. 1993. Heat flow from the Earth's interior: analysis of the global data set. *Rev. Geophys.* 31(3):267–80
- Pons ML, Fujii T, Rosing M, Quitté G, Télouk P, Albarède F. 2013. A Zn isotope perspective on the rise of continents. *Geobiology* 11(3):201–14
- Pujol M, Marty B, Burgess R, Turner G, Philippot P. 2013. Argon isotopic composition of Archaean atmosphere probes early Earth geodynamics. *Nature* 498:87–90
- Reimink JR, Chacko T, Stern RA, Heaman LM. 2014. Earth's earliest evolved crust generated in an Icelandlike setting. Nat. Geosci. 7(7):529–33
- Reimink JR, Davies JHFL, Chacko T, Stern RA, Heaman LM, et al. 2016. No evidence for Hadean continental crust within Earth's oldest evolved rock unit. *Nat. Geosci.* 9(10):777–80
- Rey PF, Coltice N. 2008. Neoarchean lithospheric strengthening and the coupling of Earth's geochemical reservoirs. *Geology* 36(8):635–38
- Rino S, Komiya T, Windley BF, Katayama I, Motoki A, Hirata T. 2004. Major episodic increases of continental crustal growth determined from zircon ages of river sands; implications for mantle overturns in the Early Precambrian. *Phys. Earth Planet. Inter.* 146:369–94
- Roberts NMW, Spencer CJ. 2014. The zircon archive of continent formation through time. Geol. Soc. Lond. Spec. Publ. 389:197–225
- Rudnick RL. 1995. Making continental crust. Nature 378(6557):571-78
- Rudnick RL, Gao S. 2003. Composition of the continental crust. In *Treatise on Geochemistry*, Vol. 3: *The Crust*, ed. RL Rudnick, pp. 1–64. Amsterdam: Elsevier
- Sandiford M, Kranendonk MJV, Bodorkos S. 2004. Conductive incubation and the origin of dome-and-keel structure in Archean granite-greenstone terrains: a model based on the eastern Pilbara Craton, Western Australia. *Tectonics* 23(1):TC1009
- Schoene B, Dudas FOL, Bowring SA, de Wit M. 2009. Sm-Nd isotopic mapping of lithospheric growth and stabilization in the eastern Kaapvaal craton. *Terra Nova* 21(3):219–28
- Scholl DW, von Huene R. 2007. Exploring the implications for continental basement tectonics if estimated rates of crustal removal (recycling) at Cenozoic subduction zones are applied to Phanerozoic and Precambrian convergent ocean margins. In 4-D Framework of Continental Crust: Memoir 200, ed. RD Hatcher Jr., MP Carlson, JH McBride, JRM Catalán, pp. 9–32. Boulder, CO: Geol. Soc. Am.
- Scholl DW, von Huene R. 2009. Implications of estimated magmatic additions and recycling losses at the subduction zones of accretionary (non-collisional) and collisional (suturing) orogens. *Geol. Soc. Lond. Spec. Publ.* 318:105–25
- Shapiro NM, Ritzwoller MH. 2002. Monte-Carlo inversion for a global shear-velocity model of the crust and upper mantle. *Geophys. J. Int.* 151(1):88–105
- Shields GA. 2007. A normalised seawater strontium isotope curve: possible implications for Neoproterozoic-Cambrian weathering rates and the further oxygenation of the Earth. *eEarth* 2:35–42
- Shirey SB, Richardson SH. 2011. Start of the Wilson cycle at 3 Ga shown by diamonds from subcontinental mantle. *Science* 333(6041):434–36
- Simon NSC, Carlson RW, Pearson DG, Davies GR. 2007. The origin and evolution of the Kaapvaal cratonic lithospheric mantle. J. Petrol. 48(3):589–625
- Sizova E, Gerya T, Brown M. 2014. Contrasting styles of Phanerozoic and Precambrian continental collision. Gondwana Res. 25(2):522–45
- Sizova E, Gerya T, Brown M, Perchuk LL. 2010. Subduction styles in the Precambrian: insight from numerical experiments. *Lithos* 116(3–4):209–29

- Sizova E, Gerya T, Stüwe K, Brown M. 2015. Generation of felsic crust in the Archean: a geodynamic modeling perspective. *Precambrian Res.* 271:198–224
- Smithies RH, Van Kranendonk MJ, Champion DC. 2007. The Mesoarchean emergence of modern-style subduction. *Gondwana Res.* 11(1–2):50–68
- Spencer CJ, Cawood PA, Hawkesworth CJ, Raub TD, Prave AR, Roberts NMW. 2014. Proterozoic onset of crustal reworking and collisional tectonics: reappraisal of the zircon oxygen isotope record. *Geology* 42(5):451–54
- Stern CR. 2011. Subduction erosion: rates, mechanisms, and its role in arc magmatism and the evolution of the continental crust and mantle. *Gondwana Res.* 20(2–3):284–308
- Stockwell CW. 1961. Structural provinces, orogenies and time classification of rocks of the Canadian Precambrian Shield. Geol. Surv. Can. Pap. 61–17:108–18, Ottawa, Can.
- Tang M, Chen K, Rudnick RL. 2016. Archean upper crust transition from mafic to felsic marks the onset of plate tectonics. *Science* 351:372–75
- Tappe S, Smart KA, Pearson DG, Steenfelt A, Simonetti A. 2011. Craton formation in Late Archean subduction zones revealed by first Greenland eclogites. *Geology* 39(12):1103–6
- Taylor SR, McLennan SM. 1985. The Continental Crust: Its Composition and Evolution: An Examination of the Geochemical Record Preserved in Sedimentary Rocks. Oxford, UK: Blackwell Sci.
- Taylor SR, McLennan SM. 1995. The geochemical evolution of the continental crust. *Rev. Geophys.* 33(2):241–65
- Turner S, Rushmer T, Reagan M, Moyen J-F. 2014. Heading down early on? Start of subduction on Earth. Geology 42(2):139–42
- Valley JW, Lackey JS, Cavosie AJ, Clechenko CC, Spicuzza MJ, et al. 2005. 4.4 billion years of crustal maturation: oxygen isotope ratios of magmatic zircon. *Contrib. Mineral. Petrol.* 150(6):561–80
- Valley JW, Peck WH, King EM, Wilde SA. 2002. A cool early Earth. Geology 30(4):351-54
- van Hunen J, van den Berg AP. 2008. Plate tectonics on the early Earth: limitations imposed by strength and buoyancy of subducted lithosphere. *Lithos* 103(1–2):217–35
- Van Kranendonk MJ, Philippot P, Lepot K, Bodorkos S, Pirajno F. 2008. Geological setting of Earth's oldest fossils in the ca. 3.5 Ga Dresser Formation, Pilbara Craton, Western Australia. *Precambrian Res.* 167(1– 2):93–124
- Van Kranendonk MJ, Smithies RH, Hickman AH, Champion DC. 2007. Secular tectonic evolution of Archean continental crust: interplay between horizontal and vertical processes in the formation of the Pilbara Craton, Australia. *Terra Nova* 19(1):1–38
- Vervoort JD, Kemp AIS. 2016. Clarifying the zircon Hf isotope record of crust-mantle evolution. *Chem. Geol.* 425:65–75
- Voice PJ, Kowalewski M, Eriksson KA. 2011. Quantifying the timing and rate of crustal evolution: global compilation of radiometrically dated detrital zircon grains. *7. Geol.* 119(2):109–26
- Walker RJ, Carlson RW, Shirey SB, Boyd FR. 1989. Os, Sr, Nd, and Pb isotope systematics of southern African peridotite xenoliths: implications for the chemical evolution of subcontinental mantle. *Geochim. Cosmochim. Acta* 53(7):1583–95
- Walter MJ. 1998. Melting of garnet peridotite and the origin of komatiite and depleted lithosphere. *J. Petrol.* 39(1):29–60
- Watts AB, Zhong SJ, Hunter J. 2013. The behavior of the lithosphere on seismic to geologic timescales. *Annu. Rev. Earth Planet. Sci.* 41(1):443–68
- Wilde SA, Valley JW, Peck WH, Graham CM. 2001. Evidence from detrital zircons for the existence of continental crust and oceans on the Earth 4.4 Gyr ago. *Lett. Nature* 409(6817):175–78
- Willbold M, Hegner E, Stracke A, Rocholl A. 2009. Continental geochemical signatures in dacites from Iceland and implications for models of early Archaean crust formation. *Earth Planet. Sci. Lett.* 279(1–2):44–52
- Worsley TR, Nance RD, Moody JB. 1986. Tectonic cycles and the history of the Earth's biogeochemical and paleoceanographic record. *Paleoceanography* 1(3):233–63
- Wu F-Y, Walker RJ, Ren X-W, Sun D-Y, Zhou X-H. 2003. Osmium isotopic constraints on the age of lithospheric mantle beneath northeastern China. *Chem. Geol.* 196(1–4):107–29
- Wu F-Y, Walker RJ, Yang Y-H, Yuan H-L, Yang J-H. 2006. The chemical-temporal evolution of lithospheric mantle underlying the North China Craton. *Geochim. Cosmochim. Acta* 70(19):5013–34

- Xu Y-G, Blusztajn J, Ma J-L, Suzuki K, Liu JF, Hart SR. 2008. Late Archean to Early Proterozoic lithospheric mantle beneath the western North China craton: Sr-Nd-Os isotopes of peridotite xenoliths from Yangyuan and Fansi. *Lithos* 102(1–2):25–42
- Young ED, Kohl IE, Warren PH, Rubie DC, Jacobson SA, Morbidelli A. 2016. Oxygen isotopic evidence for vigorous mixing during the Moon-forming giant impact. Science 351(6272):493–96
- Zhang H-F, Goldstein SL, Zhou X-H, Sun M, Cai Y. 2009. Comprehensive refertilization of lithospheric mantle beneath the North China Craton: further Os-Sr-Nd isotopic constraints. J. Geol. Soc. Lond. 166(2):249–59
- Zhang Y-L, Liu C-Z, Ge W-C, Wu F-Y, Chu Z-Y. 2011. Ancient sub-continental lithospheric mantle (SCLM) beneath the eastern part of the Central Asian orogenic belt (CAOB): implications for crust-mantle decoupling. *Lithus* 126(3–4):233–47