

*Annual Review of Cancer Biology*

# Engineering T Cells to Treat Cancer: The Convergence of Immuno-Oncology and Synthetic Biology

Joseph H. Choe,\* Jasper Z. Williams,\*  
and Wendell A. Lim

Department of Cellular and Molecular Pharmacology and Cell Design Initiative, University of California, San Francisco, California 94158, USA; email: [wendell.lim@ucsf.edu](mailto:wendell.lim@ucsf.edu)

ANNUAL  
REVIEWS **CONNECT**

[www.annualreviews.org](http://www.annualreviews.org)

- Download figures
- Navigate cited references
- Keyword search
- Explore related articles
- Share via email or social media

Annu. Rev. Cancer Biol. 2020. 4:121–39

The *Annual Review of Cancer Biology* is online at [cancerbio.annualreviews.org](http://cancerbio.annualreviews.org)

<https://doi.org/10.1146/annurev-cancerbio-030419-033657>

Copyright © 2020 by Annual Reviews. This work is licensed under a Creative Commons Attribution 4.0 International License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited. See credit lines of images or other third-party material in this article for license information

\*These authors contributed equally to this article

## Keywords

chimeric antigen receptors, cellular immunotherapy, synthetic biology, engineered immune cells, T cells, solid tumors

## Abstract

T cells engineered to recognize and kill tumor cells have emerged as powerful agents for combating cancer. Nonetheless, our ability to engineer T cells remains relatively primitive. Aside from CAR T cells for treating B cell malignancies, most T cell therapies are risky, toxic, and often ineffective, especially those that target solid cancers. To fulfill the promise of cell-based therapies, we must transform cell engineering into a systematic and predictable science by applying the principles and tools of synthetic biology. Synthetic biology uses a hierarchical approach—assembling sets of modular molecular parts that can be combined into larger circuits and systems that perform defined target tasks. We outline the toolkit of synthetic modules that are needed to overcome the challenges of solid cancers, progress in building these components, and how these modules could be used to reliably engineer more effective and precise T cell therapies.

## INTRODUCTION

### Challenges in Engineering T Cells to Treat Cancer

#### Chimeric antigen receptor (CAR):


synthetic receptor protein that combines an antigen-binding domain with a T cell-activating intracellular signaling domain; it is typically expressed on T cells to enable them to specifically recognize and kill cancer cells

#### Antigen:

a protein, peptide, or polysaccharide that can be specifically bound by antibodies or cell receptors

T cells modified to express chimeric antigen receptors (CARs), which redirect cytotoxicity toward tumor cells (**Figure 1a**), have proven to be remarkably effective for treating B cell malignancies. Such treatments demonstrate high rates of response (70–90%) in clinical trials and have resulted in the first two FDA (Food and Drug Administration)-approved genetically modified cell therapies (Bouchkouj et al. 2018, O’Leary et al. 2018, Park et al. 2016). Nonetheless, there has yet to be a clear success in engineering T cells to treat solid tumors (Klebanoff et al. 2016, Newick et al. 2016), which comprise ~90% of all cancer cases (Brown 2000). For this emerging platform to fulfill its potential, key challenges must be overcome to enhance the reliability, efficacy, and safety of T cell therapies.

CAR T cells targeting solid tumors have failed to mount effective and precise responses due to several major challenges. First, there appears to be a lack of truly cancer-specific antigens expressed by solid tumor cells, most of which are derived from and share antigen expression with healthy epithelial cells. Thus, targeting solid tumor-associated antigens has often resulted in severe and sometimes fatal on-target, off-tumor killing of normal tissues that may express the antigen, albeit at lower levels (Johnson et al. 2009; Linette et al. 2013; Morgan et al. 2010, 2013; Parkhurst et al. 2011). Second, T cells can be ineffective in immunosuppressive tumor microenvironments (TMEs) of many solid cancers (Binnewies et al. 2018). Third, there is a dearth of ways to control



**Figure 1**

CAR T cells and challenges facing them for solid tumors. (a) Diagram of CAR T structure, which includes an extracellular recognition domain (scFv) bound to the cancer antigen and fused to intracellular TCR signaling domains (CD3ζ) and costimulatory domains (e.g., CD28 or 4-1BB). (b) There are three major challenges for CAR T cells when treating solid tumors. CAR T cells must precisely recognize solid tumors while preventing normal tissue cross-reaction, overcome the suppressive tumor microenvironment, and have user control and safety platforms. Abbreviations: CAR, chimeric antigen receptor; scFv, single-chain variable fragment; TCR, T cell receptor.

the T cells after they are transferred to the patient, presenting a strong safety concern that limits our ability to test and develop more potent T cells. Here we describe ongoing efforts to advance T cell engineering that hopefully will soon enable the engineering of cells capable of executing the herculean combination of tasks necessary for effective treatment of solid cancers (Fischbach et al. 2013, Geering & Fussenegger 2015, Lim & June 2017).

## Immunotherapy Meets Synthetic Biology: Assembling a Toolkit to Systematically Program Therapeutic Cells That Can Go the Distance

A major emerging theme in therapeutic T cell engineering is the application of synthetic biology principles (Chakravarti & Wong 2015, Chen & Chen 2019, Roybal & Lim 2017, Wu et al. 2015b). The field of synthetic biology tries to understand cells as modular regulatory systems by investigating how cells are wired to give specific sense-response behaviors and, more importantly, how to re-program cells to perform novel functions (Cheng & Lu 2012, Kitada et al. 2018). Synthetic biology uses molecular parts to hierarchically assemble cellular devices and systems that perform complex tasks. While synthetic biology was originally largely focused on engineering microbes (bacteria and yeast) (Cameron et al. 2014), in the last five years there has been an explosion of mammalian cell applications. Cells, especially T cells, are an ideal chassis for therapeutic engineering due to their ability to execute more intricate behaviors than traditional small-molecule or biologic drugs. Using synthetic biology, scientists can in principle develop a toolkit of individual components that can be integrated into cellular circuits that hone the therapeutic potential of T cells.

In this review, we discuss a variety of synthetic biology modules for engineered anticancer T cells. Engineering T cells combining various individual therapeutic modules into cohesive circuits will be necessary to eliminate solid cancers. We propose that synthetic biology efforts to engineer T cells should be driven by three major needs: (a) enhancing the tumor recognition precision to prevent healthy tissue cross-reaction/toxicity, (b) boosting the ability to overcome suppressive TMEs, and (c) enabling user control over engineered cells to enhance safety in patients (**Figure 1b**). The broader application of engineered T cells in patient care also faces other issues such as cell source [allogeneic versus autologous (Qasim et al. 2017, Yang et al. 2015)], manufacturing (Esensten et al. 2016, Vormittag et al. 2018), and cost (Sarkar et al. 2018). These issues are critical for the future of engineered T cell therapies but are beyond the scope of this review.

## SMARTER RECOGNITION OF CANCER

### Challenges of Solid Tumor Recognition: Balancing Precision with Flexibility

FDA-approved CAR T cells recognize a single B cell lineage antigen (CD19) and potently kill both malignant and normal B cells. While anti-CD19 CAR T cells are not truly cancer specific, their on-target, off-tumor effects cause manageable toxicities, as B cells are relatively expendable (Brudno & Kochenderfer 2016, Kochenderfer et al. 2012). However, elimination of normal tissues sharing the CAR antigen usually cannot be tolerated. In fact, there have been several instances of lethal cross-reaction, where CAR T cells targeting tumor antigens have cross-reacted with normal tissues expressing low antigen levels, demonstrating the need for T cells that detect cancer more precisely (Klebanoff et al. 2016, Rosenberg & Restifo 2015). Another tumor recognition challenge is that most cancers have heterogeneous antigen expression (Gerlinger et al. 2012; McGranahan & Swanton 2015, 2017; Sigalotti et al. 2004). While certain antigens may be expressed by many cancer cells within a tumor, often there are tumor cells with no or low levels of antigen, which can lead to the development of tumor resistance. Thus, CAR T cells targeting a single antigen

---


**On-target/off-tumor toxicity:** a side effect of CAR T cell therapy in which the engineered T cells attack normal cells that express the antigen being targeted by the CAR T cell

**Tumor microenvironment (TME):**

the environment surrounding the tumor, which is influenced by the tumor to promote tumor growth and immune suppression

**Synthetic biology:** an approach to redesign existing biological systems through the design and construction of new biological entities such as genetic circuits, synthetic proteins, and signaling pathways

---



**Figure 2**

Antigen-sensing strategies to improve tumor-targeting precision. (a) The emerging synthetic biology toolkit for cell engineering allows for multiple-antigen-sensing T cells, which enable smart sense-and-response approaches leading to more precise T cell activation and killing. (b) CCR-CAR T cells are transduced with CARs that provide limited activation upon binding of one antigen. Only when both CAR and CCR antigens are bound are the T cells fully activated, enabling AND gate logic. (c) synNotch-CAR AND gates allow for robust combinatorial antigen gating due to sequential activation steps. An engineered T cell is initially unprimed and does not express a CAR until the synNotch receptor binds to a given antigen, A. Once primed by antigen A, the CAR is expressed and the T cell can be activated and kill any cancer cells expressing another antigen, B. (d) CARs cannot be activated when iCARs engage with a specific normal tissue antigen even if CARs are engaging with its cognate antigen, allowing for NOT gate logic. (e) Conventional CARs (green) indiscriminately kill both normal cells expressing low levels of antigen and cancer cells expressing high levels of antigen. Density-sensing circuit CARs (purple) can discriminate between low and high levels of antigen and can therefore eliminate only the cancer cells. Abbreviations: CAR, chimeric antigen receptor; CCR, chimeric costimulatory receptor; ECM, extracellular matrix; iCAR, inhibitory CAR; synNotch, synthetic Notch; TF, transcription factor.

lack the flexibility to capture heterogeneous antigen expression patterns in solid tumors (Chen et al. 2018). To improve targeting specificity and flexibility, researchers are using synthetic biology approaches to program T cells to recognize combinations of antigens, sense antigen density, and target heterogeneous antigens (Figure 2a) (Ebert et al. 2018).

### Combinatorial Antigen Recognition Can Improve Tumor Targeting Precision

One approach to improve tumor targeting precision is to engineer T cells with Boolean AND gate recognition—T cells that only kill in response to sensing two antigens on cancer cells and

spare healthy cells expressing only one antigen. An early AND gate strategy drew inspiration from the two-signal model for T cell activation (Chen & Flies 2013). The primary signal occurs when the T cell receptor (TCR) binds its cognate antigen; however, the primary signal by itself is insufficient to enable full T cell activation and proliferative response. A secondary signal from a costimulatory receptor is required to achieve full activation. FDA-approved CAR T cells combine both primary (CD3 $\zeta$ ) and costimulatory (CD28 or 4-1BB) signaling within one receptor. The primary and costimulatory signals can be split between two receptors, each targeting a different antigen, creating CAR T cells with an AND logic gate (**Figure 2b**) (Kloss et al. 2013, Wilkie et al. 2012). In this strategy, T cells are cotransduced with (a) a CAR with a low-affinity single-chain variable fragment (scFv) targeting antigen A and the CD3 $\zeta$  signaling domain and (b) a chimeric costimulatory receptor (CCR) with an scFv targeting antigen B and a costimulatory signaling domain. T cells engineered with a CAR and a CCR could kill target cells that expressed just the CAR antigen *in vitro*, but after scFv affinity tuning, both antigens were required to achieve optimal clearance of mouse xenograft tumors (Kloss et al. 2013).

Recently, our lab developed a new class of receptors called synthetic Notch (synNotch) receptors that can be used to build even more robust AND gates (Morsut et al. 2016; Roybal et al. 2016a,b). synNotch receptors (based on the native Notch receptor) use an extracellular recognition domain (e.g., scFv) to detect a target antigen. Binding the target triggers a proteolytic cleavage event that releases the intracellular domain of the receptor. In the synNotch receptor, this domain is a synthetic transcription factor that when released can enter the nucleus and drive expression of user-specified transgenes linked to the responsive promoter. Thus, synNotch circuits require transduction of the receptor and the response promoter. We have shown that T cells engineered with synNotch-driven CAR expression can function as highly precise and robust AND gates, sparing single-antigen but killing dual-antigen tumors in preclinical mouse models (**Figure 2c**). The synNotch-CAR AND gate strategy has shown significantly greater *in vitro* killing specificity for dual-antigen cancer cells than the CAR-CCR strategy because the synNotch-CAR mechanism of AND gating requires that the T cells transit through a series of sequential cell states (unprimed, primed, and activated), making activation by single-antigen cells very limited. The ability to precisely discriminate single-antigen cells could be very important when the CAR antigen is also expressed by a highly sensitive organ, such as the brain or lung. More recently, because of their flexibility and modularity, we have shown that synNotch receptors can be harnessed in combination with CARs and TCRs to generate a wide variety of different Boolean gates for diverse combinatorial antigen-sensing applications (W.A. Lim, unpublished data).

NOT gates are important yet underdeveloped antigen recognition modules that inhibit T cells from killing cross-reactive normal cells. T cells engineered with A-AND-NOT-B circuits are activated by antigen A but are dominantly inhibited by antigen B. Employing NOT gates in combinatorial antigen recognition circuits should help prevent toxicities to healthy tissues known to also express the CAR antigen. The only published NOT gate method for engineered T cells relies on expression of an inhibitory CAR (iCAR) that fuses an extracellular antigen-specific scFv to an intracellular inhibitory signaling domain [e.g., PD-1 (programmed cell death protein 1) or CTLA-4 (cytotoxic T lymphocyte-associated antigen 4)] (**Figure 2d**) (Fedorov et al. 2013). Coexpression of an iCAR with a CAR can prevent T cell killing of target cells that express the iCAR antigen. iCAR inhibitory effects are temporary and reversible, which is desirable for T cell NOT gates. However, iCARs' ability to inhibit T cell killing is highly dependent on high receptor and antigen expression levels and the CAR signaling architecture. Thus, we need to develop more robust and versatile T cell NOT gates, as bioinformatic analysis of antigen expression patterns in a variety of cancers and healthy tissues suggests that negative regulation will be the most discriminatory component in combinatorial antigen-recognition circuits (O. Troyanskaya, personal communication).

---

### T cell receptor

**(TCR):** composed of  $\alpha\beta$  subunits displaying immunoglobulin-like variable domains that recognize peptide antigens associated with major histocompatibility complex (MHC) molecules critical for T cell development, activation, and differentiation

**Logic gate:** integrated circuits that are used to implement Boolean logic operations such as AND, OR, and NOT circuits

### Single-chain variable fragment (scFv):

synthetic protein derived from an antibody that fuses the variable regions of the heavy and light chains of immunoglobulins, which can be used to give antigen specificity to a receptor or protein

### Synthetic Notch (SynNotch) receptor:

synthetic receptor protein that combines an antigen-binding domain with a transcriptional activator domain that can drive expression of user-specified target genes

---

## Sensing Antigen Density as a Mechanism to Discriminate Between Cancer and Normal Cells

Despite the safety of antibody therapy targeting ERBB2, a CAR built using this antibody, Herceptin® (trastuzumab), caused fatal toxicity due to targeting normal cells expressing low levels of antigen (Morgan et al. 2010). It is now known that CAR T cells can target antigens expressed at significantly lower levels than traditional mAb (monoclonal antibody) therapies, increasing the risk of toxicity (Stone et al. 2012, Walker et al. 2017, Watanabe et al. 2018). One approach to boost the specificity of cancer recognition is to incorporate antigen density sensing, as tumors often overexpress antigens found at lower levels in normal tissues. Density sensing was first demonstrated by lowering CAR scFv affinity to only recognize antigen-overexpressing cells (Caruso et al. 2015, Liu et al. 2015). An affinity-tuned T cell should robustly kill tumor cells overexpressing the antigen and spare normal cells expressing physiologic levels (**Figure 2e**). To achieve affinity-tuned CAR recognition of a wide pool of antigens, researchers will likely need to carry out specific screens to generate lower-affinity antibodies than are generated using conventional methods (Lim & June 2017).

Our lab is exploring alternative methods to engineer antigen density sensing in T cells. New circuits that incorporate antibody affinity, receptor expression levels, and positive feedback loops can achieve even higher discrimination based on antigen density than by tuning CAR scFv affinity alone (W.A. Lim, unpublished data). Moving forward, combinatorial antigen recognition circuits could also be improved by integrating density-sensing capabilities.

## Overcoming Antigen Loss or Heterogeneity by Cotargeting Multiple Antigens for Killing

Another disadvantage of single-antigen-targeting CARs is that cancer cells within tumors often heterogeneously express antigens or downregulate target antigen expression during treatment, both of which can cause escape. For example, cancers can lose expression of CD19 via a variety of genetic resistance mechanisms, and a disappointing proportion of patients who initially respond to CD19 CAR T cells relapse due to antigen loss (Bagashev et al. 2018, Orlando et al. 2018, Sotillo et al. 2015). These relapses sparked the development of a variety of systems to program T cells with OR gates that kill based on recognition of either CD19 or CD22 (or another B cell antigen) (Zah et al. 2016, Zhao et al. 2019). Perhaps the most promising strategy is to engineer T cells with one CAR that has tandem antigen recognition domains separately targeting CD19 and CD22 (Fry et al. 2017, Schultz et al. 2018, Qin et al. 2018). OR gate CARs can also be built using a single binding domain that can bind multiple tumor antigens, which is usually achieved using a natural ligand as the targeting domain (Baumeister et al. 2018, Gilham & Maher 2017, Klampatsa et al. 2017, Lee et al. 2017). Other approaches include expressing up to three separate CARs (or a CAR and a TCR) targeting different antigens in the same T cell or dosing multiple T cells, each targeting a different antigen (Bielamowicz et al. 2017, Chen et al. 2017, Ruella et al. 2016, Slaney et al. 2017). However, expressing one CAR with tandem recognition domains has shown the best efficacy in preclinical mouse models (Hegde et al. 2016).

Engineering T cells to overcome antigen heterogeneity is critical when targeting solid tumors. For example, in glioblastoma multiforme, a subset of patients express a truly tumor-specific antigen, deletion variant III of the epidermal growth factor receptor (EGFRvIII); however, EGFRvIII is heterogeneously expressed, and a CAR T cell clinical trial targeting EGFRvIII failed to observe any tumor regression despite evidence of killing EGFRvIII-positive cancer cells (O'Rourke et al. 2017). Thus, generating CAR T cell systems that are specific enough to prevent toxicities but also flexible enough to detect heterogeneous antigen expression patterns will be critical for treating




solid tumors. We are currently exploring how synNotch-CAR circuits could be used to prime based on recognition of a heterogeneous antigen like EGFRvIII, but then kill based on a more homogeneous, though less specific, antigen. Overall, approaches that allow T cells to integrate information across multiple cells in a tumor would be powerful tools in overcoming heterogeneity, and could be another unique advantage of cell therapies.

## OVERCOMING THE SUPPRESSIVE TUMOR MICROENVIRONMENT

### Multiple Suppressive Mechanisms Limit T Cell Activity Within Solid Tumors

Solving the recognition problem to safely target cancer cells is only one piece of the puzzle for CAR T cells. Solid tumors limit anti-tumor T cell activity by using multiple immunosuppressive factors such as inhibitory cells (e.g., regulatory T cells), soluble signals [e.g., TGF- $\beta$  (transforming growth factor beta)], cell-cell contact ligands (e.g., PD-1 axis), and physical barriers [extracellular matrix (ECM)] (**Figure 3**). TMEs are also characterized by hypoxia and nutrient deprivation that metabolically inhibit T cell cytotoxicity (Anderson et al. 2017, Binnewies et al. 2018, Jerby-Arnon et al. 2018). CAR T cell clinical trials for solid tumors have produced disappointing results; therefore, we need better ways to engineer T cells to traffic to, infiltrate into, and overcome suppressive signals in TMEs. Synthetic biology can help develop a versatile toolkit of modularly combinable



**Figure 3**

Engineering T cells to overcome inhibitory tumor microenvironment. T cells engineered with the conventional CARs are susceptible to tumor immune suppression via checkpoints/activation-induced cell death ligands, inhibitory cytokines, and physical barriers (ECM). Conventional CAR T cells can be improved by equipping T cells with synthetic tools such as ECM-degrading enzymes, deletion of the PD-1 or Fas genes, and expression of dominant-negative cytokine receptors and additional synthetic receptors enabling tumor-inducible responses. The next generation of CAR T cells can actively remodel the tumor microenvironment and become resistant to tumor suppression. Abbreviations: CAR, chimeric antigen receptor; ECM, extracellular matrix.

parts to engineer disease-specific T cells with customizable circuits to overcome the unique suppressive mechanisms presented by different tumor types.

---

**Chemokines:**

chemotactic cytokines that direct migration/positioning of immune cells within the body

**Immune checkpoint:**

inhibitory and stimulatory pathways that are critical for self-tolerance and regulating immune response; cancer cells often inhibit antitumor immune response using the immune checkpoint pathway

---

## Improving T Cell Trafficking to and Infiltration into Tumors

CAR T cell therapy for solid tumors is limited by trafficking and infiltration (Idorn & Thor Straten 2018, Slaney et al. 2014). Since blood cancers circulate in the same compartments as CAR T cells, engineered tumor homing has not been necessary. However, solid cancers are outside of the lymphoid circulation; therefore, synthetically homing T cells to solid tumors could help. Making matters more difficult, tumors often secrete chemokines that prevent T cell homing (Harlin et al. 2009), and T cells often have incompatible chemokine receptors for chemokines found in tumors (Griffith et al. 2014). To enhance tumor homing, groups have cotransduced CAR T cells with chemokine receptors corresponding to chemokines found in particular tumors. This approach has been tested in preclinical mouse models of Hodgkin's lymphoma, mesothelioma, and neuroblastoma (Craddock et al. 2010, Kershaw et al. 2002, Moon et al. 2011, Stasi et al. 2009).

Rather than relying on endogenous chemokine/chemokine receptor pairs, our group has shown that fully synthetic systems can control cell motility. Using a G protein-coupled receptor modified to respond to a bioinert drug, we engineered immune cells with drug-directed migration (Park et al. 2014). This technology can be combined with a drug-releasing bead implanted at a disease site to direct only engineered cells to migrate toward the drug. Still, we need better methods to control cell trafficking, and new strategies to enable recruitment of tunable quantities of engineered cells to tumors would significantly push the field forward.

Solid tumors also have physical barriers that pose additional challenges to T cells. The ECM in solid tumors limits T cells' penetration and aggregation (Peranzoni et al. 2013, Salmon et al. 2012). Heparan sulfate proteoglycan (HSPG) is an important tumor ECM component. By engineering CAR T cells to express an HSPG-degrading enzyme, a group achieved increased infiltration and antitumor activity in preclinical tumor models (**Figure 3**) (Caruana et al. 2015). However, without some form of local regulation, T cells with ECM-degrading payloads may cause toxicity.

## Overcoming Immune Checkpoint and Cytokine Inhibition

Another class of mechanisms that cancers use to inhibit immunity is expression of immunosuppressive ligands or cytokines (Iwai et al. 2002, Leach et al. 1996, Mariathasan et al. 2018, Rabinovich et al. 2007). These signals can critically regulate T cell fate/function and decrease antitumor efficacy. One approach to address this hurdle is to reduce T cells' inhibitory signaling capabilities. CAR T cells with PD-1 expression disrupted by Cas9-mediated gene editing (Rupp et al. 2017) or TALEN (transcription activator-like effector nuclease) editing (Menger et al. 2016) have enhanced the clearance of PD-L1-expressing tumors (**Figure 3**). Alternatively, CAR T cells engineered to constitutively secrete mAbs/scFvs blocking the PD-1 axis led to local accumulation of the immune checkpoint inhibitors within mouse xenograft tumors and enhanced clearance (Rafiq et al. 2018, Suarez et al. 2016).

Another approach to overcome suppressive signals is to express dominant-negative receptors that act as ligand sinks and block signaling. TGF- $\beta$  is a cytokine secreted by tumors that can cause apoptotic effects (Li et al. 2006); however, tumors can avoid the apoptotic effects by expressing a nonfunctional TGF- $\beta$  receptor (Knaus et al. 1996, Park et al. 1994). Overexpressing a signaling-incompetent TGF- $\beta$  receptor in CAR T cells has enhanced proliferation, cytokine secretion, and persistence in preclinical models (**Figure 3**) (Bollard et al. 2002, Kloss et al. 2018). Another approach to overcome suppression is to synthetically rewire a naturally inhibitory



input to a stimulatory output. For example, expression of a chimeric molecule that swaps the cytoplasmic tail of the CD200R inhibitory receptor with that of the costimulatory receptor CD28 caused enhanced T cell proliferation and effector function in preclinical models (Kretz-Rommel et al. 2007, Oda et al. 2017). Other similar receptors that block inhibitory signals or transduce them into stimulatory signals have also been developed to help CAR T cells overcome TMEs (Figure 3) (Liu et al. 2016, Yamamoto et al. 2019). While these are promising ways to boost efficacy, interfering with these inhibitory checkpoint and cytokine pathways in engineered T cells will require tight regulation to avoid activating autoimmunity (June et al. 2017).

### Overcoming Metabolic Suppression: New Checkpoints on the Block

Due to high nutrient consumption rates by cancers cells, tumors create suppressive metabolic microenvironments that inhibit immunity (Pearce et al. 2013, Wang & Green 2012). For example, the glycolytic metabolite phosphoenolpyruvate (PEP) sustains NFAT (nuclear factor of activated T cells) signaling and effector function, and insufficient PEP levels cause diminished antitumor T cell responses. Overexpressing PEP carboxykinase 1, which catalyzes PEP production, in T cells led to greater effector function (Ho et al. 2015). These results show that modulating T cell metabolism can enhance function.

Given nutrient deficits in solid tumors, CAR T cells' metabolic characteristics are critical for effective therapy. CAR costimulatory domain identity is known to influence the T cell metabolic state. For example, CD28 CAR costimulation drives T cell differentiation into effector memory cells with enhanced glycolysis, while 4-1BB costimulation promotes central memory T cell formation with increased respiratory capacity, increased fatty acid oxidation, and enhanced mitochondrial biogenesis (Kawalekar et al. 2016). CAR T cell studies have shown that CD28 signaling leads to early proliferation but decreased persistence, while 4-1BB signaling leads to increased persistence and lasting memory formation (Long et al. 2015, Zhang et al. 2007). There is a clear need for improved CAR costimulatory signaling to give CAR T cells the most metabolically advantageous system to balance proliferation and persistence.

### Promoting T Cell Memory, Proliferation, and Persistence

Methods to induce memory CAR T cells that balance proliferation and persistence would improve the efficacy of T cell therapies for both hematological and solid cancers. A new approach to solve this problem is to reduce the strength or duration of CAR signaling. One study showed that using CARs with a single immunoreceptor tyrosine-based activation motif (ITAM), rather than the three ITAMs normally found in CD3 $\zeta$ , led to highly functional T cells with enhanced persistence in preclinical tumor models (Feucht et al. 2019). The reduced activation signal from the CAR with a single ITAM caused especially profound increases in efficacy relative to traditional CARs in experiments infusing very low T cell numbers.

Another approach is to regulate T cells' epigenetic profile to maintain a proliferative state long-term. One study reported that anti-CD19 CAR transgene insertion unexpectedly disrupted the epigenetic regulator *TET2* gene in one T cell during the lentiviral vector-mediated integration (Fraietta et al. 2018). The *TET2* gene disruption caused this single T cell to proliferate massively in the patient, and at the peak of response, 94% of CAR T cells were found to have expanded from that single clone. Demonstrating the ability to intentionally engineer this behavior, the group knocked down the *TET2* gene and found that CAR T cell proliferation, persistence, and potency-enhancing effects were recapitulated. Similar effects on CAR function have been achieved by expressing altered c-Jun proteins in engineered T cells to drive optimal proliferative

phenotypes (Lynn et al. 2019). While potentially promising, modifications that enhance T cell proliferation and persistence create the risk of causing T cells to become autoreactive or cancerous. Thus, it may be important in the future not to simply make constitutive changes to the T cells to increase their potency (i.e., constitutive expression or knockout of particular genes), but rather to make these changes inducible at the site of the tumor (**Figure 3**). Smarter T cells that can autonomously sense when and where to amplify their potency and durability may be ideal.

## **USER CONTROL AND SAFETY**

### **The Need to Regulate Modifications that Enhance T Cell Activity to Prevent Dangerous, Out-of-Control Immune Activation**

Enhancing the potency of engineered T cells has been a major priority; however, if not carefully controlled, this could cause serious side effects. We currently lack robust ways to regulate engineered T cells once they are in patients. Enhanced safety measures, including methods for user control, will be critical to enable widespread adoption of CAR T cell therapy. Many methods are in development to control the strength, duration, and location of CAR T cell activity. Ultimately, the goal is to engineer T cells that can autonomously regulate themselves to prevent adverse events. However, even in the hypothetical context of self-regulating engineered T cells, interventional means to control cell activity during treatment are still desirable.


### **Gene Editing Strategies to Improve Predictability and Safety of Engineered T Cells**

The current standard approach engineers CAR T cells using viral vectors that permanently integrate transgenes in the genome (Esensten et al. 2016, Levine et al. 2017). Virally transduced CAR T cells experience random integration, making them less predictable and reliable. This may result in unintended consequences—such as clonal expansion, oncogenic transformation, and variable expression of the transgene—that could be avoided if the transgene were integrated into a specific locus. One study directed the CAR transgene into the TRAC (T cell receptor  $\alpha$  constant) locus using CRISPR/Cas9 genome editing technology and found more uniform CAR expression with enhanced antitumor effects at lower doses of T cells, which could reduce side effects associated with infusing high T cell numbers (Eyquem et al. 2017).

Gene editing has also been used to inactivate toxicity-related genes in CAR T cells. Cytokine release syndrome (CRS) is a major CAR T cell toxicity (Brentjens et al. 2013, Brudno & Kochenderfer 2016). Although CRS can be managed with anti-IL-6R $\alpha$  monoclonal antibodies or glucocorticoids (Le et al. 2018, Maude et al. 2014a), it can still be life threatening (Porter et al. 2018). To prevent CRS, a group inactivated the granulocyte macrophage colony-stimulating factor (GM-CSF) gene, which has been known to promote CRS, in CAR T cells using TALENs (Sachdeva et al. 2019). Upon activation, engineered CAR T cells with inactivated GM-CSF induced decreased macrophage-dependent secretion of CRS biomarkers without impairing T cell cytotoxicity or proliferation.

### **Drug-Controlled Synthetic Systems to Control T Cell Activity for Increased Safety**

Many safety and control systems in development use small molecules or biologics to regulate the location, intensity, or duration engineered T cell activity. The earliest type of control system was



**Figure 4**

Methods to control T cell activity and population size during treatment to enhance safety. (a) A strategy for constructing a kill switch CAR (chimeric antigen receptor) expressing an inducible caspase 9 with a drug-binding domain. When the small-molecule dimerizer is present, it initiates a signaling cascade leading to apoptosis of the engineered cell. (b) Truncated epidermal growth factor receptor (tEGFR) can be expressed on T cells to be used as a kill switch. Monoclonal antibodies against the EGFR can be utilized to mediate antibody-dependent cellular cytotoxicity or complement-dependent cytotoxicity and cause CAR T cells to lyse. (c) A strategy for constructing an ON switch CAR includes splitting the key CAR components from the conventional CAR into two separate polypeptides that can be conditionally brought together when a heterodimerizing small-molecule agent is present. (d) Unlike the conventional CAR T cell activation, which is mediated by direct recognition of a cancer antigen, adaptor-mediated CARs engage with tumor cells by an adaptor, which includes a peptide neoepitope (PNE) specifically recognized by the anti-PNE CAR and a cancer antigen-specific small-chain variable fragment (scFv).

kill switches to eliminate CAR T cells in case of severe toxicity (**Figure 4a**) (Berger et al. 2006, Bonini et al. 1997). A popular kill switch for drug-induced CAR T cell apoptosis (iCasp9) was developed by fusing a modified human caspase 9 to the FK506 binding protein (FKBP) that dimerizes in the presence of a small-molecule drug (Straathof et al. 2005). An advantage of this system is that the drug eliminates only engineered T cells and ignores native immune cells, and it has shown efficacy in both preclinical and clinical contexts (Diaconu et al. 2017, Stasi et al. 2011). Furthermore, FDA-approved small molecules such as rapamycin can control iCasp9 switches with alternative heterodimerization domains and may face fewer regulatory hurdles (Stavrou et al. 2018).

Systems have also been developed that use antibodies to eliminate engineered T cells. Compact epitope tag constructs with low probability of natural immunogenicity were built for binding by FDA-approved antibodies. FDA-approved antibodies rituximab and cetuximab have been used with CAR T cells expressing the corresponding truncated antigen epitope constructs for both ex vivo cell selection and in vivo cell depletion and tracking (**Figure 4b**) (Philip et al. 2014,

Valton et al. 2018, Wang et al. 2011). However, relying on antibody-dependent cellular toxicity to eliminate CAR T cells is risky in cancer patients with reduced immunity due to preconditioning or other previous chemotherapy.

Rather than killing engineered T cells, another approach is to make CAR T cell activation dependent on a drug signal. Our group has developed ON switch CARs that need to sense both an antigen and a small molecule to activate T cells (**Figure 4c**) (Wu et al. 2015a). The ON switch CAR separates the key intracellular signaling components from the extracellular antigen binding domain on separate polypeptides that contain partner drug-inducible heterodimerization domains. The ON switch CAR is inactive until it senses both heterodimerizing drug and antigen. This system allows T cell activity to be titratably and reversibly inhibited.

Another group developed a system in which CD3 $\zeta$  signaling is initiated by antigen binding while costimulatory signaling is independently controlled by a small-molecule dimerizer (Foster et al. 2017). T cells with this system are designed to fully activate only when both the target-specific CD3 $\zeta$  signal and the small-molecule inducible costimulatory signal are received; however, like the CCR AND gate system, in vitro and some in vivo killing of CD3 $\zeta$  CAR antigen-positive cells occur in the absence of the dimerizer drug. Still, the same group showed that this drug-dependent costimulatory system could be used in the same CAR T cells as an orthogonal drug-dependent inducible caspase-9-based safety switch (Duong et al. 2018). Robust, independent control of both CAR T cell expansion/persistence and death using orthogonal drug switches could enable strict modulation of therapeutic T cell population size throughout treatment.


### **Adaptor-Mediated CARs Combine Drug-Controlled Safety with Ability to Combat Resistance by Switching Antigen Targeting**

FDA-approved CAR T cells lack activity control after antigen binding and do not allow for retargeting of different antigens without using a new CAR T cell product. To address these limitations, researchers have split signaling and antigen-binding components between separate molecules in adaptor-mediated CARs (Cartellieri et al. 2016, Kudo et al. 2014, Ma et al. 2016). For example, in the peptide neoepitope (PNE) CAR system, T cells are engineered to express a CAR targeting a short, bio-orthogonal peptide tag. Under normal conditions, the PNE CAR does not bind target tumors (**Figure 4d**) (Rodgers et al. 2016, Viaud et al. 2018). When a tumor antigen-specific antibody fragment fused to the PNE tag is introduced, anti-PNE CAR T cells are redirected to kill the tumors cells. This system allows for reversible titration of CAR activity without the need to eliminate the CAR T cell. Furthermore, the same engineered T cells can change targets by adding a different tag-fused antibody.

Recently, a group developed a similar split, universal, and programmable (SUPRA) CAR (Cho et al. 2018). The SUPRA CAR system is a two-component receptor system composed of various pairs of universal orthogonal receptors expressed by T cells and corresponding tumor-targeting scFv adaptors that engage the receptors through leucine zipper interactions. Antigen targeting can be swapped by changing the adaptor, and signaling outcome is determined by the identity of the universal receptor. Like other adaptor-mediated CARs, the SUPRA CAR allows for reversible and titratable CAR activity, with the added ability to send combinations of signals to T cells for functions such as combinatorial antigen sensing.

## **CONCLUSION AND OUTLOOK**

Adoptive T cell therapy for cancer is at an exciting but critical point in its development. The success and FDA approval of anti-CD19 CAR T cells for treatment of B cell cancers has clearly



**Figure 5**

Vision for next-generation therapeutic T cell engineering for solid tumors. The next generation of therapeutic T cells will need to be designed to precisely and effectively recognize cancer and not normal tissues, they must be able to overcome the suppressive microenvironment of the tumor, and users must be able to control their activity. Precise tumor biology analysis with a synthetic immunology toolkit has the potential to provide T cells capable of overcoming solid tumors.

demonstrated the powerful potential of using engineered immune cells to eliminate cancers. However, this technology is at a primitive stage, like aviation in the pioneering days of the Wright brothers. While it was clear that powered flight was possible, it was a risky, life-threatening, and unreliable endeavor—a long way from the reliability and safety of modern commercial aviation. Similarly, today's cell therapies are highly risky and unpredictable. Most attempts to engineer T cells to treat solid tumors have led to either ineffective outcomes or highly toxic cross-reactivity.

We want the power of synthetic biology to be harnessed to develop a new foundational platform for engineering more powerful, yet safer T cell therapies. We postulate that ideal cell therapies for attacking solid cancers must fulfill multiple functions: They must be designed to precisely recognize the cancer and not critical normal tissues, they must be able to overcome the suppressive TMEs, and we must be able to control their activity (**Figure 5**). Using synthetic biology, we should be able to generate more tools to program T cells capable of sensing and responding to various diseases in increasingly autonomous and regulated manners. Synthetic biology provides unique opportunities to overcome challenges identified with the first wave of T cell therapies. Synthetic biology principles and technology platforms should facilitate the broader scientific community's ability to engineer improved, innovative T cell therapies.

## DISCLOSURE STATEMENT

The authors are not aware of any affiliations, memberships, funding, or financial holdings that might be perceived as affecting the objectivity of this review.

## LITERATURE CITED

Anderson KG, Stromnes IM, Greenberg PD. 2017. Obstacles posed by the tumor microenvironment to T cell activity: a case for synergistic therapies. *Cancer Cell* 31(3):311–25

- Bagashev A, Sotillo E, Tang C-H, Black KL, Perazzelli J, et al. 2018. CD19 Alterations emerging after CD19-directed immunotherapy cause retention of the misfolded protein in the endoplasmic reticulum. *Mol. Cell. Biol.* 38(21):e00383-18
- Baumeister SH, Murad J, Werner L, Daley H, Trebeden-Negre H, et al. 2018. Phase 1 trial of autologous CAR T cells targeting NKG2D ligands in patients with AML/MDS and multiple myeloma. *Cancer Immunol. Res.* 7(1):100-12
- Berger C, Flowers ME, Warren EH, Riddell SR. 2006. Analysis of transgene-specific immune responses that limit the in vivo persistence of adoptively transferred HSV-TK-modified donor T cells after allogeneic hematopoietic cell transplantation. *Blood* 107(6):2294-302
- Bielamowicz K, Fousek K, Byrd TT, Samaha H, Mukherjee M, et al. 2017. Trivalent CAR T-cells overcome interpatient antigenic variability in glioblastoma. *Neuro-Oncology* 20(4):506-18
- Binnewies M, Roberts EW, Kersten K, Chan V, Fearon DF, et al. 2018. Understanding the tumor immune microenvironment (TIME) for effective therapy. *Nat. Med.* 24(5):541-50
- Bollard CM, Rössig C, Calonge JM, Huls HM, Wagner H-J, et al. 2002. Adapting a transforming growth factor  $\beta$ -related tumor protection strategy to enhance antitumor immunity. *Blood* 99(9):3179-87
- Bonini C, Ferrari G, Verzeletti S, Servida P, Zappone E, et al. 1997. HSV-TK gene transfer into donor lymphocytes for control of allogeneic graft-versus-leukemia. *Science* 276(5319):1719-24
- Bouchkouj N, Kasamon YL, de Claro AR, George B, Lin X, et al. 2018. FDA approval summary: axicabtagene ciloleucel for relapsed or refractory large B-cell lymphoma. *Clin. Cancer Res.* 25(6):1702-8
- Brentjens RJ, Davila ML, Riviere I, Park J, Wang X, et al. 2013. CD19-targeted T cells rapidly induce molecular remissions in adults with chemotherapy-refractory acute lymphoblastic leukemia. *Sci. Transl. Med.* 5(177):177ra38
- Brown JM. 2000. Exploiting the hypoxic cancer cell: mechanisms and therapeutic strategies. *Mol. Med. Today* 6(4):157-62
- Brudno JN, Kochenderfer JN. 2016. Toxicities of chimeric antigen receptor T cells: recognition and management. *Blood* 127(26):3321-30
- Cameron ED, Bashor CJ, Collins JJ. 2014. A brief history of synthetic biology. *Nat. Rev. Microbiol.* 12(5):381-90
- Cartellieri M, Feldmann A, Koristka S, Arndt C, Löff S, et al. 2016. Switching CAR T cells on and off: a novel modular platform for retargeting of T cells to AML blasts. *Blood Cancer J.* 6(8):e458
- Caruana I, Savoldo B, Hoyos V, Weber G, Liu H, et al. 2015. Heparanase promotes tumor infiltration and antitumor activity of CAR-redirected T lymphocytes. *Nat. Med.* 21(5):524-29
- Caruso HG, Hurton LV, Najjar A, Rushworth D, Ang S, et al. 2015. Tuning sensitivity of CAR to EGFR density limits recognition of normal tissue while maintaining potent antitumor activity. *Cancer Res.* 75(17):3505-18
- Chakravarti D, Wong WW. 2015. Synthetic biology in cell-based cancer immunotherapy. *Trends Biotechnol.* 33(8):449-61
- Chen K, Wada M, Pinz K, Liu H, Shuai X, et al. 2017. A compound chimeric antigen receptor strategy for targeting multiple myeloma. *Leukemia* 32(2):402-12
- Chen L, Flies DB. 2013. Molecular mechanisms of T cell co-stimulation and co-inhibition. *Nat. Rev. Immunol.* 13(4):227-42
- Chen LC, Chen YY. 2019. Outsmarting and outmuscling cancer cells with synthetic and systems immunology. *Curr. Opin. Biotechnol.* 60:111-18
- Chen N, Li X, Chintala NK, Tano ZE, Adusumilli PS. 2018. Driving CARs on the uneven road of antigen heterogeneity in solid tumors. *Curr. Opin. Immunol.* 51:103-10
- Cheng AA, Lu TK. 2012. Synthetic biology: an emerging engineering discipline. *Annu. Rev. Biomed. Eng.* 14:155-78
- Cho J, Collins JJ, Wong WW. 2018. Universal chimeric antigen receptors for multiplexed and logical control of T cell responses. *Cell* 173(6):1426-38.e11
- Craddock JA, Lu A, Bear A, Pule M, Brenner MK, et al. 2010. Enhanced tumor trafficking of GD2 chimeric antigen receptor T cells by expression of the chemokine receptor CCR2b. *J. Immunother.* 33(8):780-88



- Diaconu I, Ballard B, Zhang M, Chen Y, West J, et al. 2017. Inducible caspase-9 selectively modulates the toxicities of CD19-specific chimeric antigen receptor-modified T cells. *Mol. Ther.* 25:580–92
- Duong MT, Collinson-Pautz MR, Morschl E, Lu A, Szymanski SP, et al. 2018. Two-dimensional regulation of CAR-T therapy with orthogonal switches. *Mol. Ther. Oncolyt.* 12:124–37
- Ebert LM, Yu W, Gargett T, Brown MP. 2018. Logic-gated approaches to extend the utility of chimeric antigen receptor T-cell technology. *Biochem. Soc. Trans.* 46(2):391–401
- Esensten JH, Bluestone JA, Lim WA. 2016. Engineering therapeutic T cells: from synthetic biology to clinical trials. *Annu. Rev. Pathol. Mech. Dis.* 12:305–30
- Eyquem J, Mansilla-Soto J, Giavridis T, van der Stegen SJ, Hamieh M, et al. 2017. Targeting a CAR to the TRAC locus with CRISPR/Cas9 enhances tumour rejection. *Nature* 543(7643):113–17
- Fedorov VD, Themeli M, Sadelain M. 2013. PD-1- and CTLA-4-based inhibitory chimeric antigen receptors (iCARs) divert off-target immunotherapy responses. *Sci. Transl. Med.* 5(215):215ra172
- Feucht J, Sun J, Eyquem J, Ho Y-J, Zhao Z, et al. 2019. Calibration of CAR activation potential directs alternative T cell fates and therapeutic potency. *Nat. Med.* 25(1):82–88
- Fischbach MA, Bluestone JA, Lim WA. 2013. Cell-based therapeutics: the next pillar of medicine. *Sci. Transl. Med.* 5(179):179ps7
- Foster AE, Mahendravada A, Shinnars NP, Chang W-C, Crisostomo J, et al. 2017. Regulated expansion and survival of chimeric antigen receptor-modified T cells using small molecule-dependent inducible MyD88/CD40. *Mol. Ther.* 25:2176–88
- Fraietta JA, Nobles CL, Sammons MA, Lundh S, Carty SA, et al. 2018. Disruption of TET2 promotes the therapeutic efficacy of CD19-targeted T cells. *Nature* 558(7709):307–12
- Fry TJ, Shah NN, Orentas RJ, Stetler-Stevenson M, Yuan CM, et al. 2017. CD22-targeted CAR T cells induce remission in B-ALL that is naive or resistant to CD19-targeted CAR immunotherapy. *Nat. Med.* 24(1):20–28
- Geering B, Fussenegger M. 2015. Synthetic immunology: modulating the human immune system. *Trends Biotechnol.* 33(2):65–79
- Gerlinger M, Rowan AJ, Horswell S, Larkin J, Endesfelder D, et al. 2012. Intratumor heterogeneity and branched evolution revealed by multiregion sequencing. *N. Engl. J. Med.* 366(10):883–92
- Gilham DE, Maher J. 2017. ‘Atypical’ CAR T cells: NKG2D and Erb-B as examples of natural receptor/ligands to target recalcitrant solid tumors. *Immunotherapy* 9(9):723–33
- Griffith JW, Sokol CL, Luster AD. 2014. Chemokines and chemokine receptors: positioning cells for host defense and immunity. *Annu. Rev. Immunol.* 32:659–702
- Harlin H, Meng Y, Peterson AC, Zha Y, Tretiakova M, et al. 2009. Chemokine expression in melanoma metastases associated with CD8<sup>+</sup> T-cell recruitment. *Cancer Res.* 69(7):3077–85
- Hegde M, Mukherjee M, Grada Z, Pignata A, Landi D, et al. 2016. Tandem CAR T cells targeting HER2 and IL13Rα2 mitigate tumor antigen escape. *J. Clin. Invest.* 126(8):3036–52
- Ho P-C, Bihuniak J, Macintyre AN, Staron M, Liu X, et al. 2015. Phosphoenolpyruvate is a metabolic checkpoint of anti-tumor T cell responses. *Cell* 162(6):1217–28
- Idorn M, Thor Straten P. 2018. Chemokine receptors and exercise to tackle the inadequacy of T cell homing to the tumor site. *Cells* 7(8):108
- Iwai Y, Ishida M, Tanaka Y, Okazaki T, Honjo T, Minato N. 2002. Involvement of PD-L1 on tumor cells in the escape from host immune system and tumor immunotherapy by PD-L1 blockade. *PNAS* 99(19):12293–97
- Jerby-Arnon L, Shah P, Cuoco MS, Rodman C, Su M-J, et al. 2018. A cancer cell program promotes T cell exclusion and resistance to checkpoint blockade. *Cell* 175(4):984–97.e24
- Johnson LA, Morgan RA, Dudley ME, Cassard L, Yang JC, et al. 2009. Gene therapy with human and mouse T-cell receptors mediates cancer regression and targets normal tissues expressing cognate antigen. *Blood* 114(3):535–46
- June CH, Warshauer JT, Bluestone JA. 2017. Is autoimmunity the Achilles’ heel of cancer immunotherapy? *Nat. Med.* 23(5):540–47
- Kawalekar OU, O’Connor RS, Fraietta JA, Guo L, McGettigan SE, et al. 2016. Distinct signaling of coreceptors regulates specific metabolism pathways and impacts memory development in CAR T cells. *Immunity* 44(2):380–90

- Kershaw MH, Wang G, Westwood JA, Pachynski RK, Tiffany LH, et al. 2002. Redirecting migration of T cells to chemokine secreted from tumors by genetic modification with CXCR2. *Hum. Gene Ther.* 13(16):1971–80
- Kitada T, DiAndreth B, Teague B, Weiss R. 2018. Programming gene and engineered-cell therapies with synthetic biology. *Science* 359(6376):eaad1067
- Klampatsa A, Achkova DY, Davies DM, Parente-Pereira AC, Woodman N, et al. 2017. Intracavitary ‘T4 immunotherapy’ of malignant mesothelioma using pan-ErbB re-targeted CAR T-cells. *Cancer Lett.* 393:52–59
- Klebanoff CA, Rosenberg SA, Restifo NP. 2016. Prospects for gene-engineered T cell immunotherapy for solid cancers. *Nat. Med.* 22(1):26–36
- Kloss CC, Condomines M, Cartellieri M, Bachmann M, Sadelain M. 2013. Combinatorial antigen recognition with balanced signaling promotes selective tumor eradication by engineered T cells. *Nat. Biotechnol.* 31(1):71–75
- Kloss CC, Lee J, Zhang A, Chen F, Melenhorst J, et al. 2018. Dominant negative TGF- $\beta$  receptor enhances PSMA targeted human CAR T cell proliferation and augments tumor eradication in prostate cancer. *Mol. Ther.* 26:1855–66
- Knaus P, Lindemann D, DeCoteau J, Perlman R, Yankelev H, et al. 1996. A dominant inhibitory mutant of the type II transforming growth factor beta receptor in the malignant progression of a cutaneous T-cell lymphoma. *Mol. Cell. Biol.* 16(7):3480–89
- Kochenderfer JN, Dudley ME, Feldman SA, Wilson WH, Spaner DE, et al. 2012. B-cell depletion and remissions of malignancy along with cytokine-associated toxicity in a clinical trial of anti-CD19 chimeric-antigen-receptor-transduced T cells. *Blood* 119(12):2709–20
- Kretz-Rommel A, Qin F, Dakappagari N, Ravey PE, McWhirter J, et al. 2007. CD200 expression on tumor cells suppresses antitumor immunity: new approaches to cancer immunotherapy. *J. Immunol.* 178(9):595–605
- Kudo K, Imai C, Lorenzini P, Kamiya T, Kono K, et al. 2014. T lymphocytes expressing a CD16 signaling receptor exert antibody-dependent cancer cell killing. *Cancer Res.* 74(1):93–103
- Le RQ, Li L, Yuan W, Shord SS, Nie L, et al. 2018. FDA approval summary: tocilizumab for treatment of chimeric antigen receptor T cell-induced severe or life-threatening cytokine release syndrome. *Oncology* 23(8):943–47
- Leach DR, Krummel MF, Allison JP. 1996. Enhancement of antitumor immunity by CTLA-4 blockade. *Science* 271(5256):1734–36
- Lee L, Draper B, Chaplin N, Philip B, Chin M, et al. 2017. An APRIL based chimeric antigen receptor for dual targeting of BCMA and TACI in multiple myeloma. *Blood* 131(7):746–58
- Levine BL, Miskin J, Wonnacott K, Keir C. 2017. Global manufacturing of CAR T cell therapy. *Mol. Ther. Methods Clin. Dev.* 4:92–101
- Li MO, Wan YY, Sanjabi S, Robertson A-KL, Flavell RA. 2006. Transforming growth factor- $\beta$  regulation of immune responses. *Annu. Rev. Immunol.* 24:99–146
- Lim WA, June CH. 2017. The principles of engineering immune cells to treat cancer. *Cell* 168(4):724–40
- Linette GP, Stadtmauer EA, Maus MV, Rapoport AP, Levine BL, et al. 2013. Cardiovascular toxicity and titin cross-reactivity of affinity-enhanced T cells in myeloma and melanoma. *Blood* 122(6):863–71
- Liu X, Jiang S, Fang C, Yang S, Olalere D, et al. 2015. Affinity-tuned ErbB2 or EGFR chimeric antigen receptor T cells exhibit an increased therapeutic index against tumors in mice. *Cancer Res.* 75(17):3596–607
- Liu X, Ranganathan R, Jiang S, Fang C, Sun J, et al. 2016. A chimeric switch-receptor targeting PD1 augments the efficacy of second-generation CAR T cells in advanced solid tumors. *Cancer Res.* 76(6):1578–90
- Long AH, Haso WM, Shern JF, Wanhainen KM, Murgai M, et al. 2015. 4-1BB costimulation ameliorates T cell exhaustion induced by tonic signaling of chimeric antigen receptors. *Nat. Med.* 21(6):581–90
- Lynn RC, Weber EW, Gennert D, Sotillo E, Xu P, et al. 2019. C-Jun overexpressing CAR-T cells are exhaustion-resistant and mediate enhanced antitumor activity. bioRxiv 653725. <https://doi.org/10.1101/653725>

- Ma JS, Kim J, Kazane SA, Choi S, Yun H, et al. 2016. Versatile strategy for controlling the specificity and activity of engineered T cells. *PNAS* 113(4):E450–58
- Mariathasan S, Turley SJ, Nickles D, Castiglioni A, Yuen K, et al. 2018. TGF $\beta$  attenuates tumour response to PD-L1 blockade by contributing to exclusion of T cells. *Nature* 554(7693):544–48
- Maude SL, Barrett D, Teachey DT, Grupp SA. 2014a. Managing cytokine release syndrome associated with novel T cell-engaging therapies. *Cancer J.* 20(2):119–22
- McGranahan N, Swanton C. 2015. Biological and therapeutic impact of intratumor heterogeneity in cancer evolution. *Cancer Cell* 27(1):15–26
- McGranahan N, Swanton C. 2017. Clonal heterogeneity and tumor evolution: past, present, and the future. *Cell* 168(4):613–28
- Menger L, Sledzinska A, Bergerhoff K, Vargas F, Smith J, et al. 2016. TALEN-mediated inactivation of PD-1 in tumor-reactive lymphocytes promotes intratumoral T-cell persistence and rejection of established tumors. *Cancer Res.* 76(8):2087–93
- Moon EK, Carpenito C, Sun J, Wang L-CS, Kapoor V, et al. 2011. Expression of a functional CCR2 receptor enhances tumor localization and tumor eradication by retargeted human T cells expressing a mesothelin-specific chimeric antibody receptor. *Clin. Cancer Res.* 17(14):4719–30
- Morgan RA, Chinnasamy N, Abate-Daga D, Gros A, Robbins PF, et al. 2013. Cancer regression and neurological toxicity following anti-MAGE-A3 TCR gene therapy. *J. Immunother.* 36(2):133–51
- Morgan RA, Yang JC, Kitano M, Dudley ME, Laurencot CM, Rosenberg SA. 2010. Case report of a serious adverse event following the administration of T cells transduced with a chimeric antigen receptor recognizing ERBB2. *Mol. Ther.* 18(4):843–51
- Morsut L, Roybal KT, Xiong X, Gordley RM, Coyle SM, et al. 2016. Engineering customized cell sensing and response behaviors using synthetic Notch receptors. *Cell* 164(4):780–91
- Newick K, O'Brien S, Moon E, Albelda SM. 2016. CAR T cell therapy for solid tumors. *Annu. Rev. Med.* 68:139–52
- Oda SK, Daman AW, Garcia NM, Wagener F, Schmitt T, et al. 2017. A CD200R-CD28 fusion protein appropriates an inhibitory signal to enhance T-cell function and therapy of murine leukemia. *Blood* 130(22):2410–19
- O'Leary MC, Lu X, Huang Y, Lin X, Mahmood I, et al. 2018. FDA approval summary: tisagenlecleucel for treatment of patients with relapsed or refractory B-cell precursor acute lymphoblastic leukemia. *Clin. Cancer Res.* 25(4):1142–46
- Orlando EJ, Han X, Tribouley C, Wood PA, Leary RJ, et al. 2018. Genetic mechanisms of target antigen loss in CAR19 therapy of acute lymphoblastic leukemia. *Nat. Med.* 24(10):1504–6
- O'Rourke DM, Nasrallah MP, Desai A, Melenhorst JJ, Mansfield K, et al. 2017. A single dose of peripherally infused EGFRvIII-directed CAR T cells mediates antigen loss and induces adaptive resistance in patients with recurrent glioblastoma. *Sci. Transl. Med.* 9(399):eaaa0984
- Park JH, Geyer MB, Brentjens RJ. 2016. CD19-targeted CAR T-cell therapeutics for hematologic malignancies: interpreting clinical outcomes to date. *Blood* 127(26):3312–20
- Park JS, Rhau B, Hermann A, McNally KA, Zhou C, et al. 2014. Synthetic control of mammalian-cell motility by engineering chemotaxis to an orthogonal bioinert chemical signal. *PNAS* 111(16):5896–901
- Park K, Kim S, Bang Y, Park J, Kim N, et al. 1994. Genetic changes in the transforming growth factor beta (TGF-beta) type II receptor gene in human gastric cancer cells: correlation with sensitivity to growth inhibition by TGF-beta. *PNAS* 91(19):8772–76
- Parkhurst MR, Yang JC, Langan RC, Dudley ME, Nathan D-AN, et al. 2011. T cells targeting carcinoembryonic antigen can mediate regression of metastatic colorectal cancer but induce severe transient colitis. *Mol. Ther.* 19(3):620–26
- Pearce EL, Poffenberger MC, Chang C-H, Jones RG. 2013. Fueling immunity: insights into metabolism and lymphocyte function. *Science* 342(6155):1242454
- Peranzoni E, Rivas-Caicedo A, Bougherara H, Salmon H, Donnadieu E. 2013. Positive and negative influence of the matrix architecture on antitumor immune surveillance. *Cell Mol. Life Sci.* 70(23):4431–48
- Philip B, Kokalaki E, Mekkaoui L, Thomas S, Straathof K, et al. 2014. A highly compact epitope-based marker/suicide gene for easier and safer T-cell therapy. *Blood* 124(8):1277–87

- Porter D, Frey N, Wood PA, Weng Y, Grupp SA. 2018. Grading of cytokine release syndrome associated with the CAR T cell therapy tisagenlecleucel. *J. Hematol. Oncol.* 11(1):35
- Qasim W, Zhan H, Samarasinghe S, Adams S, Amrolia P, et al. 2017. Molecular remission of infant B-ALL after infusion of universal TALEN gene-edited CAR T cells. *Sci. Transl. Med.* 9(374):eaaj2013
- Qin H, Ramakrishna S, Nguyen S, Fountaine TJ, Ponduri A, et al. 2018. Preclinical development of bivalent chimeric antigen receptors targeting both CD19 and CD22. *Mol. Ther. Oncol.* 11:127–37
- Rabinovich GA, Gabrilovich D, Sotomayor EM. 2007. Immunosuppressive strategies that are mediated by tumor cells. *Annu. Rev. Immunol.* 25:267–96
- Rafiq S, Yeku OO, Jackson HJ, Purdon TJ, van Leeuwen DG, et al. 2018. Targeted delivery of a PD-1-blocking scFv by CAR-T cells enhances anti-tumor efficacy in vivo. *Nat. Biotechnol.* 36(9):847–56
- Rodgers DT, Mazagova M, Hampton EN, Cao Y, Ramadoss NS, et al. 2016. Switch-mediated activation and retargeting of CAR-T cells for B-cell malignancies. *PNAS* 113(4):E459–68
- Rosenberg SA, Restifo NP. 2015. Adoptive cell transfer as personalized immunotherapy for human cancer. *Science* 348(6230):62–68
- Roybal KT, Lim WA. 2017. Synthetic immunology: hacking immune cells to expand their therapeutic capabilities. *Annu. Rev. Immunol.* 35:229–53
- Roybal KT, Rupp LJ, Morsut L, Walker WJ, McNally KA, et al. 2016a. Precision tumor recognition by T cells with combinatorial antigen-sensing circuits. *Cell* 164(4):770–79
- Roybal KT, Williams JZ, Morsut L, Rupp LJ, Kolinko I, et al. 2016b. Engineering T cells with customized therapeutic response programs using synthetic Notch receptors. *Cell* 167(2):419–32.e16
- Ruella M, Barrett DM, Kenderian SS, Shestova O, Hofmann TJ, et al. 2016. Dual CD19 and CD123 targeting prevents antigen-loss relapses after CD19-directed immunotherapies. *J. Clin. Invest.* 126(10):3814–26
- Rupp LJ, Schumann K, Roybal KT, Gate RE, Ye CJ, et al. 2017. CRISPR/Cas9-mediated PD-1 disruption enhances anti-tumor efficacy of human chimeric antigen receptor T cells. *Sci. Rep.* 7:737
- Sachdeva M, Duchateau P, Depil S, Poirot L, Valton J. 2019. Granulocyte-macrophage colony-stimulating factor inactivation in CAR T-cells prevents monocyte-dependent release of key cytokine release syndrome mediators. *J. Biol. Chem.* 294(14):5430–37
- Salmon H, Franciszkiewicz K, Damotte D, Dieu-Nosjean M-C, Validire P, et al. 2012. Matrix architecture defines the preferential localization and migration of T cells into the stroma of human lung tumors. *J. Clin. Invest.* 122(3):899–910
- Sarkar RR, Gloude NJ, Schiff D, Murphy JD. 2018. Cost-effectiveness of chimeric antigen receptor T-cell therapy in pediatric relapsed/refractory B-cell acute lymphoblastic leukemia. *J. Natl. Cancer Inst.* 111(7):719–26
- Schultz L, Davis KL, Baggott C, Chaudry C, Marcy A, et al. 2018. Phase 1 study of CD19/CD22 bispecific chimeric antigen receptor (CAR) therapy in children and young adults with B cell acute lymphoblastic leukemia (ALL). *Blood* 132(Suppl. 1):898
- Sigalotti L, Fratta E, Coral S, Tanzarella S, Danielli R, et al. 2004. Intratumor heterogeneity of cancer/testis antigens expression in human cutaneous melanoma is methylation-regulated and functionally reverted by 5-Aza-2'-deoxycytidine. *Cancer Res.* 64(24):9167–71
- Slaney CY, Kershaw MH, Darcy PK. 2014. Trafficking of T cells into tumors. *Cancer Res.* 74(24):7168–74
- Slaney CY, von Scheidt B, Davenport AJ, Beavis PA, Westwood JA, et al. 2017. Dual-specific chimeric antigen receptor T cells and an indirect vaccine eradicate a variety of large solid tumors in an immunocompetent, self-antigen setting. *Clin. Cancer Res.* 23(10):2478–90
- Sotillo E, Barrett DM, Black KL, Bagashev A, Oldridge D, et al. 2015. Convergence of acquired mutations and alternative splicing of CD19 enables resistance to CART-19 immunotherapy. *Cancer Discov.* 5(12):1282–95
- Stasi A, Angelis B, Rooney CM, Zhang L, Mahendravada A, et al. 2009. T lymphocytes coexpressing CCR4 and a chimeric antigen receptor targeting CD30 have improved homing and antitumor activity in a Hodgkin tumor model. *Blood* 113(25):6392–402

- Stasi A, Tey S-K, Dotti G, Fujita Y, Kennedy-Nasser A, et al. 2011. Inducible apoptosis as a safety switch for adoptive cell therapy. *N. Engl. J. Med.* 365(18):1673–83
- Stavrou M, Philip B, Traynor-White C, Davis CG, Onuoha S, et al. 2018. A rapamycin activated caspase 9 based suicide gene. *Mol. Ther.* 26:1266–76
- Stone JD, Aggen DH, Schietinger A, Schreiber H, Kranz DM. 2012. A sensitivity scale for targeting T cells with chimeric antigen receptors (CARs) and bispecific T-cell engagers (BiTEs). *Oncoimmunology* 1(6):863–73
- Straathof KC, Pulè MA, Yotnda P, Dotti G, Vanin EF, et al. 2005. An inducible caspase 9 safety switch for T-cell therapy. *Blood* 105(11):4247–54
- Suarez E, Chang D-K, Sun J, Sui J, Freeman GJ, et al. 2016. Chimeric antigen receptor T cells secreting anti-PD-L1 antibodies more effectively regress renal cell carcinoma in a humanized mouse model. *Oncotarget* 7(23):34341–55
- Valton J, Guyot V, Boldajipour B, Sommer C, Pertel T, et al. 2018. A versatile safeguard for chimeric antigen receptor T-cell immunotherapies. *Sci. Rep.* 8:8972
- Viaud S, Ma JS, Hardy IR, Hampton EN, Benish B, et al. 2018. Switchable control over in vivo CAR T expansion, B cell depletion, and induction of memory. *PNAS* 115(46):E10898–906
- Vormittag P, Gunn R, Ghorashian S, Veraitch FS. 2018. A guide to manufacturing CAR T cell therapies. *Curr. Opin. Biotechnol.* 53:164–81
- Walker AJ, Majzner RG, Zhang L, Wanhainen K, Long AH, et al. 2017. Tumor antigen and receptor densities regulate efficacy of a chimeric antigen receptor targeting anaplastic lymphoma kinase. *Mol. Ther.* 25:2189–201
- Wang R, Green DR. 2012. Metabolic checkpoints in activated T cells. *Nat. Immunol.* 13(10):907–15
- Wang X, Chang W-C, Wong CW, Colcher D, Sherman M, et al. 2011. A transgene-encoded cell surface polypeptide for selection, in vivo tracking, and ablation of engineered cells. *Blood* 118(5):1255–63
- Watanabe K, Kuramitsu S, Posey AD, June CH. 2018. Expanding the therapeutic window for CAR T cell therapy in solid tumors: the knowns and unknowns of CAR T cell biology. *Front. Immunol.* 9:2486
- Wilkie S, van Schalkwyk MC, Hobbs S, Davies DM, van der Stegen SJ, et al. 2012. Dual targeting of ErbB2 and MUC1 in breast cancer using chimeric antigen receptors engineered to provide complementary signaling. *J. Clin. Immunol.* 32(5):1059–70
- Wu C-Y, Roybal KT, Puchner EM, Onuffer J, Lim WA. 2015a. Remote control of therapeutic T cells through a small molecule-gated chimeric receptor. *Science* 350(6258):aab4077
- Wu C-Y, Rupp LJ, Roybal KT, Lim WA. 2015b. Synthetic biology approaches to engineer T cells. *Curr. Opin. Immunol.* 35:123–30
- Yamamoto TN, Lee P-H, Vodnala SK, Gurusamy D, Kishton RJ, et al. 2019. T cells genetically engineered to overcome death signaling enhance adoptive cancer immunotherapy. *J. Clin. Investig.* 129:1551–65
- Yang Y, Jacoby E, Fry TJ. 2015. Challenges and opportunities of allogeneic donor-derived CAR T cells. *Curr. Opin. Hematol.* 22(6):509–15
- Zah E, Lin M-Y, Silva-Benedict A, Jensen MC, Chen YY. 2016. T cells expressing CD19/CD20 bispecific chimeric antigen receptors prevent antigen escape by malignant B cells. *Cancer Immunol. Res.* 4(6):498–508
- Zhang H, Snyder KM, Suhoski MM, Maus MV, Kapoor V, et al. 2007. 4-1BB is superior to CD28 costimulation for generating CD8<sup>+</sup> cytotoxic lymphocytes for adoptive immunotherapy. *J. Immunol.* 179(7):4910–18
- Zhao J, Song Y, Liu D. 2019. Clinical trials of dual-target CAR T cells, donor-derived CAR T cells, and universal CAR T cells for acute lymphoid leukemia. *J. Hematol. Oncol.* 12(1):17