

# *Annual Review of Pharmacology and Toxicology* Experimental Models of SARS-CoV-2 Infection: Possible Platforms to Study COVID-19 Pathogenesis and Potential Treatments

Sareh Pandamooz,<sup>1</sup> Benjamin Jurek,<sup>2</sup>  
Carl-Philipp Meinung,<sup>3</sup> Zahra Baharvand,<sup>4</sup>  
Alireza Sahebi Shahem-abadi,<sup>5</sup> Silke Haerteis,<sup>2</sup>  
Jaleel A. Miyan,<sup>6</sup> James Downing,<sup>7</sup>  
Mehdi Dianatpour,<sup>1</sup> Afshin Borhani-Haghighi,<sup>8</sup>  
and Mohammad Saied Salehi<sup>8</sup>

<sup>1</sup>Stem Cells Technology Research Center, Shiraz University of Medical Sciences, Shiraz, Iran;  
email: Sareh.pandamooz@gmail.com

<sup>2</sup>Institute for Molecular and Cellular Anatomy, University of Regensburg, Regensburg 93053,  
Germany

<sup>3</sup>Department of Molecular and Behavioural Neurobiology, University of Regensburg,  
Regensburg 93053, Germany

<sup>4</sup>Department of Cellular and Molecular Biology, Faculty of Biological Sciences, Kharazmi  
University, Tehran, Iran

<sup>5</sup>Department of Animal Biology, Faculty of Biological Sciences, Kharazmi University, Tehran,  
Iran

<sup>6</sup>Faculty of Biology, Medicine & Health, Division of Neuroscience & Experimental Psychology,  
The University of Manchester, Manchester M13 9PL, United Kingdom

<sup>7</sup>School of Pharmacy and Biomolecular Sciences, Faculty of Science, Liverpool John Moores  
University, Liverpool L2 2QP, United Kingdom

<sup>8</sup>Clinical Neurology Research Center, Shiraz University of Medical Sciences, Shiraz, Iran;  
email: Saied\_salehi@sums.ac.ir

## ANNUAL REVIEWS **CONNECT**

[www.annualreviews.org](http://www.annualreviews.org)

- Download figures
- Navigate cited references
- Keyword search
- Explore related articles
- Share via email or social media

Annu. Rev. Pharmacol. Toxicol. 2022. 62:25–53

First published as a Review in Advance on  
February 19, 2021

The *Annual Review of Pharmacology and Toxicology* is  
online at [pharmtox.annualreviews.org](http://pharmtox.annualreviews.org)

<https://doi.org/10.1146/annurev-pharmtox-121120-012309>

Copyright © 2022 by Annual Reviews.  
All rights reserved

## Keywords

SARS-CoV-2, COVID-19, in vitro models, animal models

## Abstract

In December 2019, a novel coronavirus crossed species barriers to infect humans and was effectively transmitted from person to person, leading to a worldwide pandemic. Development of effective clinical interventions, including vaccines and antiviral drugs that could prevent or limit the


burden or transmission of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), is a global health priority. It is thus of utmost importance to assess possible therapeutic strategies against SARS-CoV-2 using experimental models that recapitulate aspects of the human disease. Here, we review available models currently being developed and used to study SARS-CoV-2 infection and highlight their application to screen potential therapeutic approaches, including repurposed antiviral drugs and vaccines. Each identified model provides a valuable insight into SARS-CoV-2 cellular tropism, replication kinetics, and cell damage that could ultimately enhance understanding of SARS-CoV-2 pathogenesis and protective immunity.

## 1. INTRODUCTION

In December 2019 in Wuhan, China, a novel coronavirus outbreak occurred that caused infectious respiratory disease. Soon after, due to its high transmissibility and pathogenicity, the virus spread across international borders. The World Health Organization (WHO) declared coronavirus disease 2019 (COVID-19) a pandemic, which has become the worst public health crisis in a century. In February 2020, the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) was identified as the causative agent of COVID-19, and as of November 10, 2020, it has infected more than 50.6 million people worldwide and caused more than 1.26 million recorded deaths. Similar to other lethal coronaviruses [i.e., severe acute respiratory syndrome coronavirus (SARS-CoV) and Middle East respiratory syndrome coronavirus (MERS-CoV)], SARS-CoV-2 triggers an excessive immune response known as a cytokine storm that can lead to multiple organ failure and death (1).

SARS-CoV-2 is a single-stranded RNA virus. Its envelope is covered with trimeric spike (S) proteins that bind to human angiotensin-converting enzyme 2 (hACE2) on the plasma membrane of host cells (2). After binding, the S protein is cleaved by the transmembrane protease serine subtype 2 (TMPRSS2), thereby facilitating viral entry (3). Following fusion of viral and cellular membranes, the virus empties its genetic contents and hijacks the translational machinery of the host cells to perpetuate itself, resulting in the production of copies of the virus at the expense of the host cells.

The main challenge in the current pandemic is the novelty of the disease, which results in unfamiliar pathophysiology and limited knowledge of SARS-CoV-2 biology (4). At the time this manuscript was submitted for publication, there were three COVID-19 vaccines for which certain national regulatory authorities had authorized the use. There are also many other potential COVID-19 vaccine candidates currently in development, and several of these are likely to be approved by the time of this publication. Nevertheless, for this pandemic and for others that will inevitably occur in the near future, it is of utmost importance to use platforms with physiologically relevant model systems that can faithfully indicate the viral life cycle, delineate the pathology, recapitulate host-virus interactions in human cells, and provide a platform to test protective treatments and therapies such as potential vaccines and antiviral drugs. Models of SARS-CoV-2 infection are also useful for evaluating other emerging strategies, for example, stem cell therapies that might combat COVID-19 (5). In this review, we summarize several model systems that have recently been used to study SARS-CoV-2 (**Figure 1**). Here, we attempt to define the hierarchy in the use of model organisms, from simple monolayer cell culture platforms to complex tissue explant cultures in Section 2. We also discuss animal models ranging from mouse models to nonhuman primates (NHPs). Please note that the authors do not include experimental models identified after November 10, 2020.



**Figure 1**

Overview of different model systems used to study severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection. A wide range of in vitro models, including primary cell culture, cell lines, induced pluripotent stem cell (iPSC)-derived stem cells, organoids, tissue explants, and organ-on-a-chip, have been used as platforms to study SARS-CoV-2 replication kinetics, cell damage profiles, and cellular tropism. Furthermore, animal models in mice, ferrets, Syrian hamsters, and nonhuman primates are used to evaluate SARS-CoV-2 infection in specific target organs. These platforms are also useful to assess systemic effects and immune responses. In addition, numerous computational and mathematical models can predict trends of coronavirus disease 2019 (COVID-19) transmission and the number of infected individuals, thereby aiding in epidemic prevention and control measures.

## 2. IN VITRO MODELS

As with SARS-CoV, MERS-CoV, and other emerging viruses, in vitro models can yield understanding of viral replication kinetics and the profile of viral-induced cell damage. Studies of SARS-CoV-2 have involved many in vitro models: primary cell culture, cell lines, induced pluripotent stem cell (iPSC)-derived cells, organ-on-a-chip, organoids, and tissue explants (**Table 1**).

Although SARS-CoV-2 primarily targets lung epithelium and causes respiratory infection, growing evidence documents that the intestine, kidney, liver, pancreas, brain, heart, blood vessels, and other organs can also be infected. Accordingly, SARS-CoV-2 infection has been assessed in numerous in vitro platforms derived from different human tissues and organs (**Table 2**).

### 2.1. Primary Cell Culture

Based on their high expression of angiotensin-converting enzyme 2 (ACE2) and TMPRSS2, human airway epithelial cells such as nasal (goblet and ciliated cells) and bronchial cells provide clinically relevant models for screening drugs against SARS-CoV-2 (6). Pizzorno et al. (7) used human airway epithelium (HAE) cultured at an air-liquid interface (ALI) to characterize viral

**Table 1 Benefits and limitations of in vitro models in COVID-19 research**

Experimental model	Benefits (references)	Limitation(s) (references)
Primary cell culture	Reproduces major structural features of human respiratory epithelia (7) Can test immune modulators and antivirals for SARS-CoV-2 infection (8)	Availability and capacity for expansion are limited Absence of immune cells (7)
Cell lines	Suitable to replicate and isolate SARS-CoV-2 (13, 14) Useful for characterizing cellular tropism, viral replication kinetics, and virus-induced cell damage profile of SARS-CoV-2 (21)	Cell line tropism might not fully represent how SARS-CoV-2 replicates and affects human organs in the physiological state (21) Constant proliferation and lack of polarization might impact the transfection efficiency of SARS-CoV-2 (24)
iPSC-derived cells	Transcriptome of iPSC-derived cells is similar to their respective primary counterparts and allows them to respond to immune stimuli (26) Useful for drug repurposing (25)	Directed differentiation of hiPSCs to desired cell types is time consuming Cellular immaturity of hiPSC-derived cells can affect their susceptibility to SARS-CoV-2 (30)
Organoids	Contain a full range of differentiated cell types as are present in the target organ Aid in understanding the tissue tropism of SARS-CoV-2 (24, 54, 57) Useful tools for antiviral drug discovery and development (55)	Typically lack vasculature and immune cells (184)
Organ-on-a-chip	Provides vascular perfusion, organ-level physical microenvironments, tissue-tissue interfaces, relevant mechanical cues, and fluid flow Human lung airway chip may be useful to expedite drug repurposing (61)	Relatively low throughput
Tissue explant	Retains cytoarchitecture and three-dimensional organization of the tissue Provides a suitable model to study tropism, replication competence, and innate immune responses of SARS-CoV-2 in the human respiratory tract (21, 68)	Ex vivo tissue explant culture is short lasting Lacks in vivo effects of host adaptive immune and systemic inflammatory responses (68) Supply of human tissue is difficult

Abbreviations: COVID-19, coronavirus disease 2019; hiPSC, human induced pluripotent stem cell; iPSC, induced pluripotent stem cell; SARS-CoV-2, severe acute respiratory syndrome coronavirus 2.

infection kinetics, tissue-level remodeling of the cellular ultrastructure, and transcriptional immune signatures induced by SARS-CoV-2. Since this model can closely recapitulate virus-host cell interactions, as in infected upper and lower airways, it has been used to evaluate the efficacy of antiviral drugs (8, 9). Primary cultures mainly characterize short-lived SARS-CoV-2 infection in HAE, but Hao et al. (10) reported data for SARS-CoV-2 in polarized HAE and ALI culture that allowed for the monitoring of SARS-CoV-2 infections for a prolonged time period. Another group used conditional cell reprogramming technology to investigate long-term cultures of cells from the human upper and lower respiratory tract (11).

## 2.2. Cell Lines

As with the use of cell culture models for characterizing cell tropism, virus-induced cell damage, and viral replication of emerging viruses, a set of cell lines have been utilized to study

**Table 2** In vitro models currently used in COVID-19 research

Model type (Origin)	Application(s)	Reference(s)
<b>Primary cell culture</b>		
Human airway epithelium	Characterizing viral infection kinetics and tissue-level remodeling of cellular ultrastructure and transcriptional immune signatures induced by SARS-CoV-2	7
	Drug screening	8
	Understanding responses of proximal and distal lung epithelium to SARS-CoV-2 infection	8, 9
	Long-term modeling of SARS-CoV-2 infection in polarized human airway epithelium	10
	Addressing SARS-CoV-2 replication	185
<b>Cell lines</b>		
VeroE6 cells, wild type (kidney epithelial cells of African green monkey)	Replicating and isolating SARS-CoV-2	3, 14, 15
VeroE6 cells, engineered to over-express TMPRSS2 or VeroE6/TMPRSS2 (kidney epithelial cells of African green monkey)	Enhanced isolation of SARS-CoV-2	13
Caco-2 cells (human epithelial colorectal adenocarcinoma)	Screening 5,632 compounds for their potential to inhibit SARS-CoV-2-induced cytotoxicity	16
	Revealing central pathways relevant for SARS-CoV-2 infection	17
Calu-3 cells (human lung adenocarcinoma)	Defining SARS-CoV-2 replication kinetics	13, 18, 20
	Drug screening (evaluating the inhibitory effect of chloroquine)	19
A549 cells (human lung adenocarcinoma)	Comparing tropism, replication kinetics, and cell damage profiling of SARS-CoV-2 and SARS-CoV	21
	Studying SARS-CoV-2 replication	15, 186
	Evaluating the expression of ACE2 and TMPRSS2	187
Huh7 cells (hepatocellular carcinoma)	Comparing tropism, replication kinetics, and cell damage profiling of SARS-CoV-2 and SARS-CoV	21
	Drug screening	193
	Defining the cellular response to the virus	188
HeLa cells (human cervical adenocarcinoma)	Determining SARS-CoV-2 cell entry mechanisms	2, 189, 194
293T cells (human embryonic kidney)	Evaluating the inhibitory activity of EK1C4 against SARS-CoV-2	190
	Assessing syncytia formation upon SARS-CoV-2 infection	191
	Testing human monoclonal antibody blockade of SARS-CoV-2 infection	192
U251 cells (human brain glioblastoma)	Comparing tropism, replication kinetics, and cell damage by SARS-CoV-2 and SARS-CoV	21
RD cells (human rhabdomyosarcoma)	Comparing tropism, replication kinetics, and cell damage by SARS-CoV-2 and SARS-CoV	21
	Evaluating the inhibitory activity of EK1C4 against SARS-CoV-2	190

(Continued)

**Table 2 (Continued)**

Model type (Origin)	Application(s)	Reference(s)
<b>Human iPSC-derived stem cells</b>		
iPSC-derived lung epithelium	Modeling SARS-CoV-2 infection and drug repurposing	25
iPSC-derived alveolar epithelial type 2 cells	Studying transcriptomic alteration after SARS-CoV-2 infection and drug screening	26
	Studying expression level of genes relevant for COVID-19 disease modeling	27
iPSC-derived airway epithelial cells	Modeling SARS-CoV-2 infection	27
iPSC-derived lung and macrophage coculture	Modeling host-pathogen interaction and immune response caused by SARS-CoV-2 infection	28
iPSC-derived cardiomyocytes	Revealing that cardiomyocytes are permissive for SARS-CoV-2 infection	29, 30
	Showing that SARS-CoV-2 infection can cause robust transcriptomic and morphological signatures of damage in cardiomyocytes	31
iPSC-derived neural progenitor cells	Assessing susceptibility of these stem cells to SARS-CoV-2 infection	34
iPSC-derived neural cells	Testing SARS-CoV-2 neurotropism	35
<b>Organoids</b>		
Lung organoids (hESC)	Identifying drug candidates that block SARS-CoV-2 entry	41
Bronchial organoids (commercially available cryopreserved human bronchial epithelial cells)	Studying SARS-CoV-2 infection and drug screening	42
Bronchial organoids (primary human bronchial epithelial cells)	Identifying potential key genes for SARS-CoV-2 infection	43
Tonsil organoids (human tonsil epithelium)	Studying SARS-CoV-2 infection	44
Intestinal organoids (human small intestinal cells)	Demonstrating that SARS-CoV-2 readily replicates in gut enterocytes	47, 48
Intestinal organoids (bat intestinal cells)	Providing evidence that SARS-CoV-2 infects bat intestinal cells	48
Intestinal organoids (human colon)	Revealing that human intestinal epithelium supports SARS-CoV-2 infection, replication, and de novo infectious virus production	49
Liver organoids (liver bile duct-derived progenitor cells)	Showing that SARS-CoV-2 infection induces cell death of cholangiocytes and bile acid accumulation	54
Liver organoids (iPSCs)	Documenting that hepatocytes and cholangiocyte organoids are permissive to SARS-CoV-2	24
Kidney organoids (human embryonic stem cells)	Revealing that human recombinant soluble ACE2 inhibits SARS-CoV-2 infection in these organoids	55
Blood vessel organoids (iPSCs)	Showing that SARS-CoV-2 infection can be reduced by human recombinant soluble ACE2	55
Brain organoids (iPSCs)	Addressing SARS-CoV-2 neurotropism Identifying that SARS-CoV-2 mainly targets neural progenitor cells and cortical neurons Evaluating therapeutic strategies	34, 35, 54, 59, 60
Brain organoids (hESCs)	Providing a model for investigating SARS-CoV-2 entry into the human brain	58
Choroid plexus organoids (iPSCs)	Showing greater SARS-CoV-2 infection of choroid plexus organoids compared to other brain regions that cause cell death and altered secretory function	35

(Continued)

Table 2 (Continued)

Model type (Origin)	Application(s)	Reference(s)
<b>Organs-On-Chips</b>		
Primary lung epithelium-on-a-chip	Modeling SARS-CoV-2 entry, replication, and host cytokine production Drug screening	62
Vascularized lung-on-a-chip	Studying SARS-CoV-2 replication	63
Microengineered alveolus-on-a-chip	Reconstituting key features of the human alveolar-capillary barrier in SARS-CoV-2 infection	64
Intestine-on-a-chip	Recapitulating intestinal injury and immune response induced by SARS-CoV-2	65
<b>Tissue explants</b>		
Human lung tissue explants	Demonstrating the viral kinetics, cell tropism, and innate immune response profile of SARS-CoV-2	68, 70
Organotypic human bronchial epithelial cells	Revealing that ciliated cells are a major target of SARS-CoV-2 infection	69, 70
Organotypic human airway epithelial cultures	Elucidating that SARS-CoV-2 causes plaque-like cytopathic effects in these cells	71
Ex vivo cultures of human conjunctiva	Showing that conjunctival epithelium is a potential portal for SARS-CoV-2 infection	70

Abbreviations: ACE2, angiotensin-converting enzyme 2; COVID-19, coronavirus disease 2019; hESC, human embryonic stem cell; iPSC, induced pluripotent stem cell; RD, rhabdomyosarcoma; SARS-CoV-2, severe acute respiratory syndrome coronavirus 2; TMPRSS2, transmembrane protease serine subtype 2.

SARS-CoV-2 infections (12). One such cell line used to replicate and isolate SARS-CoV-2 is VeroE6, originally isolated from kidney epithelial cells of an African green monkey (13–15). Caco-2, an epithelial cell line derived from a human colorectal adenocarcinoma, has been used to assess COVID-19 pathology. Ellinger et al. (16) recently used the Caco-2 cell line to screen 5,632 compounds for their potential to inhibit viral-induced cytotoxicity. Caco-2 cells have also revealed pathways relevant for SARS-CoV-2 infections, including nucleic acid metabolism, protein homeostasis, translation, and splicing (17). The Calu-3 cell line, originally derived from human lung adenocarcinoma, is another preclinical model of pulmonary viral infections that is used to study SARS-CoV-2 (13, 18–20).

Chu et al. (21) analyzed the differential susceptibility of 25 cell lines and compared the replication capability of SARS-CoV-2, as assessed by quantitative reverse transcription PCR (22). These cell lines were mainly derived from different human tissues, including from the respiratory tract (A549, Calu-3, and HFL), gastrointestinal tract (Caco-2), liver (Huh7), cervix (HeLa), kidney (293T), brain (U251), and muscle (RD). Moreover, several cell lines originated from nonhuman species such as bat (RLL, RLK, RSL, and RSK), dog (MDCK), pig (RK-15), rabbit (RK-13), cat (CRFK), porcupine (Pok), and NHPs (VeroE6, FRhK4, LLCMK2). The findings revealed that SARS-CoV-2 infected and replicated to comparable levels in human Caco-2 and Calu-3 cells over a period of 120 h.

This study provided valuable information and novel insights into SARS-CoV-2 cellular tropism, replication kinetics, and cell damage, but the authors noted several limitations with these cell lines. First, tropism among the cell lines might not exactly reflect how SARS-CoV-2 replicates and affects human organs in a physiological condition. Second, the authors used only one viral isolate (but which was highly homologous to other SARS-CoV-2 reported isolates). Third, human organs and tissues contain multiple cell types that express ACE2 heterogeneously.

Therefore, using immortalized or cancer cell lines might not fully reveal SARS-CoV-2 infection rates and mechanisms in different cell types. In addition, most of these human cancer cell lines carry tumor-associated mutations, such as in P53, which is known to regulate replication of SARS-CoV (23). This issue raises concerns as to whether these cancer cell lines recapitulate SARS-CoV-2 biology in a physiological context. Moreover, as these cells mainly originate from cancers, their proliferation and lack of polarization might impact the transfection efficiency of the virus (24).

### 2.3. iPSC-Derived Stem Cells

Although primary cell culture models have been frequently used to study SARS-CoV-2, their short in vitro maintenance is a major limitation, and because of genomic alterations, the cell lines may not faithfully respond to SARS-CoV-2 infections. By contrast, iPSCs offer the potential to study cells more reminiscent of human tissue. The transcriptome of iPSC-derived cells is similar to that of their respective primary counterparts and allows them to respond to immune stimuli. To date, several iPSC-derived cell types have provided a platform for drug development, systematic exploration of SARS-CoV-2 viral tropism, and studying cellular responses to infection.

Surendran et al. (25) generated lung epithelial cells from human iPSCs using a three-step protocol. Another research group presented an in vitro human model that could simulate the apical infection of distal lung epithelium with SARS-CoV-2 by developing iPSC-derived alveolar epithelial type 2 cells (iAT2s) as facultative progenitors of this tissue; SARS-CoV-2 infection was able to induce rapid global transcriptomic alterations in these cells (26). In addition, Abo et al. (27) compared iPSC-derived iAT2s and iPSC-derived airway epithelial cells with primary lung epithelial cell controls, concentrating on the expression levels of genes related to COVID-19 modeling. The results indicated that these two generated cells express key transcripts associated with lung identity, e.g., ACE2 and TMPRSS2. Duan et al. (28) used directed differentiation of human induced pluripotent stem cells (hiPSCs) to establish a lung/macrophage coculture system to simulate host-pathogen interaction and immune response caused by SARS-CoV-2 infection. This strategy overcomes a main concern about histocompatibility when studying human immune cells with other cell types and sustains unlimited cell resources for reliably modeling the immunology of macrophages and human lungs during infection. Using hiPSC-derived cardiomyocytes (hiPSC-CMs), Bojkova et al. (29) demonstrated that these cells are permissive for SARS-CoV-2 infection, as detected by intracellular double-stranded viral RNA and viral S protein expression. In addition to confirming the susceptibility of hiPSC-CMs to SARS-CoV-2 infection (30), it was shown that this infection can cause transcriptomic and morphological signatures of damage in these cells (31).

Accumulating evidence suggests that the human brain is another extrapulmonary target of this virus. Of note, 36% of SARS-CoV-2-infected patients displayed neurological conditions (32), and the virus has been detected in postmortem brain tissue of COVID-19 patients (33). Zhang et al. (34) reported SARS-CoV-2 infection of iPSC-derived human neural progenitor cells, while Jacob et al. (35) investigated susceptibility of iPSC-derived neurons, astrocytes, and microglia in monolayer cultures. They observed modest numbers of infected neurons and astrocytes and a high level of infection in choroid plexus epithelial cells.

### 2.4. Organoids

The emergence of human in vitro three-dimensional (3D) cell culture approaches using stem cells has received widespread attention due to their potential to overcome the limitations of classical two-dimensionally cultured and immortalized cell lines (36). Organoids are 3D structures mainly established from pluripotent stem cells (PSCs) or, alternatively, from multipotent adult stem cells resident in the tissue. They are composed of tissue-specific cell types that self-organize through

cell sorting and spatially restricted lineage commitment to generate cell assemblies with functional and architectural features of the corresponding tissue. Organoids optimally contain a full range of the differentiated cell types present in the target organ (37).

Numerous organoids have been employed to study SARS-CoV-2 infection at a level between single cells and tissue (38). As the primary SARS-CoV-2 infection target is the respiratory tract, several research groups have used lung organoids to mimic the human lung (39, 40). Han et al. (41) developed a lung organoid model using human PSCs and screened several US Food and Drug Administration (FDA)-approved drug candidates such as mycophenolic acid and imatinib as inhibitors of SARS-CoV-2 entry. Suzuki et al. (42) established human bronchial organoids (hBOs) from commercially available cryopreserved human bronchial epithelial cells. This model contained basal, club, ciliated, and goblet cell types, and treatment of SARS-CoV-2-infected hBOs with the TMPRSS2 inhibitor camostat significantly reduced viral copies in this model. Cytokine/cytokine receptor interactions, apoptosis, and P53 signaling are the main cellular mechanisms activated in these organoids following SARS-CoV-2 infections (43). Since palatine tonsils are in the front line of the immune system's defense against pathogens, tonsil epithelial cell-derived organoids reflect the distinctive characteristics of the corresponding tissue and have been proposed as a platform for studying coronavirus infections and COVID-19 biology (44).

Respiratory symptoms dominate clinical aspects of COVID-19, but involvement of the gastrointestinal system can occur as well (45, 46). A study with human small intestinal organoids suggested infection and replication of SARS-CoV-2 in human intestinal epithelium (47). Differentiated enterocytes in these organoids were readily infected by SARS-CoV-2, as demonstrated by confocal and electron microscopy. Zhou et al. (48) used expandable intestinal organoids derived from horseshoe bats of the species *Rhinolophus sinicus* that could recapitulate the bat intestinal epithelium. Due to their susceptibility to SARS-CoV-2 infections, these organoids sustain robust viral replication. The same group also confirmed SARS-CoV-2 replication in human intestinal organoids as well as upregulation of type III interferon responses. This notion was further supported by Stanifer et al. (49), who showed that SARS-CoV-2 could infect the human gastrointestinal tract and efficiently produce de novo viruses. Interestingly, they found that the type III interferon, which is naturally made by cells in response to viral infection, can be used as an antiviral strategy to protect infected cells.

The liver abundantly expresses ACE2, and liver damage can occur with SARS-CoV-2 infections (50–52). The average expression level of ACE2 in hepatocytes is 20-fold less than in cholangiocytes. This level of expression in cholangiocytes is comparable to that of lung AT2 cells (53). Zhao et al. (54) employed a previously established model of liver ductal organoids to demonstrate that these organoids are susceptible to SARS-CoV-2. RNA sequencing revealed a set of 337 differentially expressed genes in the infected organoids. The data indicate that SARS-CoV-2 infection induces cell death of cholangiocytes that can subsequently cause bile acid accumulation. This issue was further investigated by Yang et al. (24), who evaluated SARS-CoV-2 tropism in various human cells and organoids and found that human hepatocyte and cholangiocyte organoids are permissive to SARS-CoV-2 infection. Human blood vessel and kidney organoids have also been isolated, and human recombinant soluble ACE2 significantly reduced viral growth in these infected organoids (55).

Many hospitalized COVID-19 patients exhibit neurological manifestations ranging from loss of smell and headache to confusion and disabling strokes (56). Brain organoids can address several questions concerning the neurotropism and mechanisms of neuropathogenesis (57–59).

Ramani et al. (60) provided an initial insight into the unexpected neurodegenerative-like effects of SARS-CoV-2. The authors found that the Düsseldorf isolate of SARS-CoV-2 preferably targets neurons in 3D human brain organoids, leading to altered Tau distribution and phosphorylation

and eventually causing neuronal death. This finding was further supported by experiments that showed SARS-CoV-2 infection in 3D human brain organoids was localized to NESTIN-positive and TUJ1-expressing cells, suggesting that SARS-CoV-2 can directly target neural precursor cells and cortical neurons (34). Jacob et al. (35) investigated the susceptibility of region-specific brain organoids to SARS-CoV-2 and found greater infection of choroid plexus organoids that caused cell death and transcriptional dysregulation related to increased inflammation and altered secretory function. In spite of its many advantages, an important limitation of organoid technology is the lack of communication between organs as occurs in vivo. Thus, the findings require validation in more complex models. Human organoid culture remains under development, and numerous efforts to advance this technology are still in progress.

## 2.5. Organ-on-a-Chip

Organ-on-a-chip technology offers a special platform to mimic human organ infection. Human organ chips are engineered microfluidic devices populated with human cells that recapitulate tissue-tissue interfaces, fluid flows, mechanical cues, and organ-level physiology (61). Ingber and colleagues (62) developed organ chip microfluidic culture devices lined by primary human lung airway epithelium. The cells are cultured under ALI conditions and fed by continuous flow of medium. This model mimics virus entry, replication, and host cytokine production and represents a particularly useful platform to expedite drug repurposing. Recently, a vascularized lung-on-a-chip infection model was developed that consists of a coculture of primary human alveolar and human lung microvascular endothelial cells with the optional addition of CD14-positive macrophages. This model was used to study SARS-CoV-2 replication and host response to infection (63). The use of an alveolus chip was reported by Zhang et al. (64), who were able to reconstitute the key features of the alveolar-capillary barrier. Notably, they showed distinct responses of two cell types to SARS-CoV-2 infection, including activation of the type I interferon signaling cascade in epithelium and the Janus kinase–signal transducer and activator of transcription (JAK-STAT) signaling pathway in the endothelium. Guo et al. (65) created a microengineered intestine-on-a-chip device to mirror the pathophysiological features and immune responses in COVID-19. This model contains cocultured human intestinal epithelial (Caco-2) cells and mucin-secreting HT-29 cells lined in the upper channel and human umbilical vein endothelial cells in the lower channel under fluidic conditions. Collectively, organ-on-a-chip-based assays provide a platform to study SARS-CoV-2-induced pathology in real time and at high resolution in a human-relevant environment.

## 2.6. Tissue Explants

The study of human cell-cell and cell-pathogen interactions that occur in the context of tissue cytoarchitecture is important for deciphering the underlying mechanisms of many normal and pathogenic processes (66). In this regard, tissue explants or organotypic cultures are robust in vitro models that are established from surgical resection material and can be highly suitable platforms at a level between dissociated cell cultures and animal models. These models offer several unique advantages compared to other in vitro approaches: They generally retain the tissue architecture of their originated region, preserve cell-cell and cell-matrix interaction, and allow direct real-time observation (67). With respect to SARS-CoV-2 infection, human lung tissue explant is one of the most useful ex vivo models to compare the viral kinetics, cell tropism, and innate immune response profile of SARS-CoV-2 and SARS-CoV (68). This ex vivo culture showed more efficient replication but less host interferon and proinflammatory response following SARS-CoV-2

infection compared to primary cell culture. Also, single-cell RNA sequencing and electron microscopy on organotypic human bronchial epithelial cells infected with SARS-CoV-2 revealed that ciliated cells are a major target of infection (69).

Besides lung explants, ex vivo cultures of human bronchus and conjunctiva have been employed to study SARS-CoV-2 infection. These models have demonstrated that the virus can infect type 1 pneumocytes in the lung; in ciliated, mucus-secreting, and club cells of bronchial epithelium; and in conjunctival mucosa (70). In addition, Zhu et al. (71) recently characterized the replication dynamics, cell tropism, and morphogenesis of SARS-CoV-2 in organotypic human airway epithelial cultures. The results showed that SARS-CoV-2 causes plaque-like cytopathic effects in these cells. Apoptosis, cell fusion, destruction of epithelium integrity, and cilia shrinking and beaded changes were observed in the plaque-like regions. Thus, tissue explants and organotypic cultures can be useful platforms to enhance insight into SARS-CoV-2 infection.

### 3. ANIMAL MODELS FOR SARS-COV-2 RESEARCH

In vitro approaches may not provide insight into the harmful in vivo effects. There is thus an urgent need to develop animal models that are amenable to investigating effects of SARS-CoV-2 on target organs and systemic effects of the infection. A range of animal models have been developed and used (72–75) (Table 3). Each model has benefits and limitations (Table 4).

#### 3.1. Mouse

Since mouse angiotensin-converting enzyme 2 (mACE2) does not support SARS-CoV-2 binding as does hACE2, laboratory strains of mice are not readily infected by SARS-CoV-2 (76, 77). Thus, multiple interventions based on genetic modifications have been introduced to facilitate SARS-CoV-2 susceptibility in order to provide more physiologically relevant platforms for modeling the infection. In this regard, Sun et al. (78) described an adenoviral transduction-based mouse model (Ad5-hACE2-sensitized mice) that can be infected with SARS-CoV-2. These mice can develop pneumonia, characterized by weight loss, severe pulmonary pathology, and high-titer virus replication in the lungs. Thus, this model provides a tool to understand host factors involved in viral infection and clearance as well as potential therapeutic modalities. Although Bao et al. (79) recently repropounded their potentially useful hACE-2 transgenic mice for SARS-CoV-2 studies, the viral replication in these mice was suboptimal, with pathological changes in the lung and weight loss being minimal. Another suggested model is cytokeratin-18 (K18)-hACE2 transgenic mice in which hACE2 expression is driven by the epithelial cell K18 promoter. These transgenic mice share many features of severe COVID-19 infection (80). Jiang and colleagues introduced a HFH4-hACE2 transgenic mice that could be infected with SARS-CoV-2. This model was able to recapitulate a number of the symptoms and pathology seen in COVID-19 patients and pre-exposure to SARS-CoV-2 protected these mice from severe pneumonia (81). A mouse model generated using clustered regularly interspaced short palindromic repeat (CRISPR)–CRISPR-associated 9 knock-in technology to express hACE2 was predicted to be a useful platform to study SARS-CoV-2 infection and transmission, but fatalities were not observed in this model (82). Hassan et al. (83) transiently transduced a replication-defective adenovirus encoding hACE2 into the lungs of commercially available mice, which are permissive for SARS-CoV-2. This model is widely available for high-throughput drug and vaccine testing. Pujhari & Rasgon (84) developed two new strategies for generating mice with humanized lungs and immune systems as a model to define the mechanisms of SARS-CoV-2 infection. The first focused on humanized mice with ectopically grafted human lung (85), while the second concentrated on the generation of functional lungs via conditional blastocyst complementation using PSCs (86).

**Table 3 In vivo models used in COVID-19 research**

Type	Key Points	Reference(s)
Ad5-hACE2-sensitized mice	hACE2 was provided by adenovirus transduction Mouse model that can be infected with SARS-CoV-2; can aid in understanding host factors involved in viral infection/clearance and potential therapeutics	78
hACE2 transgenic mice	Mice generated by microinjection of the mouse Ace2 promoter driving the human ACE2-coding sequence into the pronuclei of fertilized ova from ICR mice Model used to study SARS-CoV-2 pathogenicity	79
K18-hACE2 transgenic mice	hACE2 expression driven by epithelial cell K18 promoter Introduced as a model to define the basis of lung disease in SARS-CoV-2 infection	80
HFH4-hACE2 transgenic mice	Mice express human ACE2 under the control of a lung ciliated epithelial cell-specific HFH4/FOXJ1 promoter Model used to recapitulate a number of infection symptoms and pathology in COVID-19 patients	81
hACE2 mice	Humanized ACE2 mouse by CRISPR-Cas9 knock-in technology A tool to study SARS-CoV-2 transmission and pathogenesis and to evaluate vaccines and therapeutics	82
hACE2 transduced mice	Mice transduced with adenoviruses encoding human ACE2 Model is now widely available for high-throughput drug and vaccine testing	83
Wild-type BALB/c mice	Uses mouse-adapted SARS-CoV-2 strain at passage 6 (called MASCP6) Model used to validate protective efficacy of candidate vaccines	87
Mouse-adapted SARS-CoV-2 (SARS-CoV-2 MA)	Uses recombinant virus that utilizes mACE2 for entry Model used to demonstrate age-related disease pathogenesis; supports the clinical use of pegylated IFN- $\lambda$ 1a as a treatment for human COVID-19	88
Syrian hamster	Infection of hamsters revealed that a SARS-CoV-2 deleted variant (Del-mut-1) has attenuated ability to cause disease in this model	89
	Surgical mask partition in hamster model can reduce SARS-CoV-2 transmission by respiratory droplets or airborne droplet nuclei	90
	Syrian hamster is a model that simulates the clinical and pathological manifestations of COVID-19; SARS-CoV-2 isolates efficiently replicate in their lungs and cause severe pathological lesions The passive transfer of convalescent serum to naïve hamsters inhibited virus replication in their lungs	92, 93
	SARS-CoV-2 is efficiently transmitted from inoculated hamsters to naïve hamsters by direct contact and via aerosols	94
	SARS-CoV-2-infected hamsters can develop neutralizing antibodies, which protect them from reinfection	92–95
	SARS-CoV-2 replication in the upper and lower respiratory tract was independent of the age of the hamster Rapid lung recovery was observed only in young hamsters	97
	Following SARS-CoV-2 infection, the STAT-dependent interferon responses play a critical role in the pathogenesis and control of the virus	98
Ferrets	Ferrets are highly susceptible to SARS-CoV-2 infection	102, 103
	Ferrets effectively transmit the virus by direct or indirect contact, recapitulating human transmission	104, 105
	The antiviral efficacies of several FDA-approved drugs against SARS-CoV-2 have been assessed in infected ferrets	106
	The alteration of gene expression from short-term to long-term SARS-CoV-2 infection was evaluated in infected ferrets	107

(Continued)

Table 3 (Continued)

Type	Key Points	Reference(s)
Nonhuman primates, general	Apes, African and Asian monkeys, and some lemurs are likely highly susceptible to SARS-CoV-2	110
Rhesus macaque	Exhibits high ACE2-spike activity and is susceptible to SARS-CoV-2 infection	111–113
	Early remdesivir treatment in SARS-CoV-2 infection had a clinical benefit	114
	SARS-CoV-2 infection induced protective immunity against subsequent reinfection	115, 116
	DNA vaccine encoding full-length S protein protected them from SARS-CoV-2 infection	117
	ChAdOx1 nCoV-19 vaccine prevented SARS-CoV-2 pneumonia	118
	Inactivated SARS-CoV-2 virus vaccine (PiCoVacc) induced SARS-CoV-2-specific neutralizing antibodies	119
	Single dose of Ad26 vector-based vaccine protected against SARS-CoV-2 infection	120
African green monkey	High level of SARS-CoV-2 replication and pronounced respiratory tract infection	121–123
Cynomolgus macaques	Useful model to study SARS-CoV-2 infection	124, 126
	NVX-CoV2373 vaccine protected against SARS-CoV-2 infection	127
Crab-eating macaques	Susceptible to SARS-CoV-2 infection	128

Abbreviations: ACE2, angiotensin-converting enzyme 2; Ad26, adenovirus serotype 26; COVID-19, coronavirus disease 2019; CRISPR-Cas9, clustered regularly interspaced short palindromic repeat–CRISPR-associated 9; FDA, US Food and Drug Administration; ICR, Institute for Cancer Research; IFN- $\lambda$ 1a, interferon  $\lambda$ 1a; K18, cytokeratin-18; SARS-CoV-2, severe acute respiratory syndrome coronavirus 2.

Table 4 Benefits and limitations of in vivo models in COVID-19 research

Experimental model	Benefits	Limitations
Mouse	Small size, ease of use, and rapid breeding make mice well suited for rapid screening of antiviral drugs and vaccines Enable exploration of effects of genetic diversity and background	Mismatched ACE2 receptors Even genetically engineered mice are not a perfect substitute for human infection
Hamster	Naturally susceptible to SARS-CoV-2 infection Infected golden Syrian hamsters closely mimic the mild to moderate disease of human patients Can be used to study viral transmission Suitable for evaluation of antiviral agents and candidate vaccines	May not reflect human pharmacokinetics May not metabolize prodrugs as do humans
Ferret	Lung and airway physiology are similar to those of humans Useful model in the study of disease transmission	May not reflect human pharmacokinetics Low viral titer in lungs
Nonhuman primate	The closest model to human infection, as their innate and adaptive immunity, physiology, and anatomy are similar to those of humans Excellent model to assess by repeated sampling and imaging, given their size and longevity Similar pharmacokinetics to humans	High costs and ethical considerations Small sample sizes

Abbreviations: ACE2, angiotensin-converting enzyme 2; COVID-19, coronavirus disease 2019; SARS-CoV-2, severe acute respiratory syndrome coronavirus 2.

Besides genetically modified mice, COVID-19 pathology has been studied in wild-type mice infected with adapted SARS-CoV-2 variants. Gu et al. (87), through inoculation with mouse-adapted SARS-CoV-2, efficiently infected young and aged BALB/c mice, resulting in moderate pneumonia and inflammatory responses. Dinno et al. (88) circumvented the inefficient interaction of viral S protein and mACE2 using reverse genetics to remodel their binding interface, resulting in a recombinant virus (SARS-CoV-2 MA) that utilizes mACE2 for entry. This SARS-CoV-2 MA replicated in both upper and lower airways of mice irrespective of their ages. Although both wild-type mouse models were suggested as reliable platforms for drug and vaccine screening, they may not fully recapitulate all aspects of the human disease (72).

### 3.2. Syrian Hamster

Syrian golden hamster is another small rodent that has been widely used to model SARS-CoV-2 infection (89, 90), although no mortality was observed after virus infection (91). Hamster ACE2 has high-affinity binding of the SARS-CoV-2 S glycoprotein receptor-binding domain. Chan et al. (92) showed that hamster ACE2 affinity to the S protein is the highest observed apart from humans and rhesus macaques. Furthermore, SARS-CoV-2 isolates efficiently replicate in hamster lungs, causing severe pathological lung lesions following intranasal infection (93). Notably, hamsters infected with SARS-CoV-2, besides showing clinical signs of disease, can transmit the virus to naïve cohoused animals by direct contact and/or via aerosols (94). Although this model can recapitulate certain aspects of human pathogenesis, infected animals can return to their original weight within 2 weeks postinfection and develop neutralizing antibodies that protect them against subsequent rechallenge with SARS-CoV-2 (95, 96). In an elegant experiment, Osterrieder et al. (97) compared the course of infection in young and aged hamsters. The authors found an identical virus replication rate in the upper and lower respiratory tract of both groups, but rapid lung recovery was only observed in young hamsters, a result of relevance to the age-dependent differences in humans. Boudewijns et al. (98), using a unique STAT2-knockout hamster, revealed a dual role of this signaling pathway in driving severe lung injury and also restricting systemic virus dissemination. This finding underlines the importance of STAT2-dependent interferon responses in the pathogenesis and control of the virus during infection and may identify new strategies for COVID-19 treatment. Suresh et al. (99) assessed the tissue-specific expression pattern of ACE2 in different organs of hamster and found that kidney, small intestine, esophagus, tongue, brain, and liver express this receptor but lung, tracheal epithelial cells, and large intestine lacked ACE2 expression. Overall, the hamster model is a useful platform to study transmission, pathogenesis, and treatment of and vaccination against SARS-CoV-2.

### 3.3. Ferret

Ferrets have been useful in reproducing human diseases caused by influenza virus and SARS-CoV, as their respiratory tract is anatomically comparable to humans and they express ACE2 abundantly on type II alveolar and granular epithelial cells in the trachea and bronchi (100, 101). In addition, ferret ACE2 contains critical SARS-CoV binding residues that render them susceptible to SARS-CoV-2 (102, 103). Infected ferrets show high virus titers in the upper respiratory tracts, and consequently, highly efficient ferret-to-ferret transmission can occur via direct or indirect contact (104). The ferret transmission pattern has been investigated by Richard et al. (105), who detected infected viruses in indirect recipients, suggesting that the SARS-CoV-2 ferret model recapitulates human-to-human transmission. Based on its susceptibility, the ferret model has been used to assess a number of FDA-approved drugs, which have been suggested as antiviral candidates against

SARS-CoV-2 and yielded results that may help facilitate the selection of antiviral treatments for COVID-19 (106). In addition, alteration of gene expression in short- and long-term-infected ferrets has been shown using bioinformatics approaches: During the course of infection, the stemness, development, and Wnt- and cytoskeleton remodeling-related pathways mainly change in the early phase of lung infection, while the extracellular matrix, immune response, G protein-coupled receptor, and cell adhesion-related pathways are affected in the later phase (107). Given the rapid spreading of SARS-CoV-2 in humans, ferrets can be a useful model for understanding COVID-19 transmission dynamics.

### 3.4. Nonhuman Primates

NHP models of SARS-CoV-2 are the gold standard for modeling human pathogenesis and for testing clinical interventions due to their genetic homology and anatomic similarity to humans (108). Based on the fidelity of NHP models for the investigation of the 2003 SARS-CoV outbreak (109), these models were reconsidered to assess SARS-CoV-2 infection. A range of NHP models have been used to study COVID-19 with varying susceptibility, which likely depends on species-specific properties of ACE2 (110). Owing to their high ACE2-spike activity, rhesus macaques are a NHP susceptible to SARS-CoV-2 infection (111–113) and have been extensively used to assess the efficacy of medical countermeasures. The efficacy of remdesivir (GS-5734), a nucleotide analog prodrug with broad antiviral activity, was investigated in a rhesus macaque model of SARS-CoV-2 (114). Remdesivir-treated animals did not show signs of respiratory disease, and pulmonary infiltration and virus load were reduced following a first dose. This study concluded that the early initiation of treatment with remdesivir can benefit infected animals. Rechallenge of infected rhesus macaques with SARS-CoV-2 resulted in a viral load reduction that suggests primary infection-induced protective immunity against reinfection in these models (115, 116).

Numerous vaccine candidates are currently being developed, and their efficacy is evaluated in NHP models. Yu et al. (117) used rhesus macaques to test a series of DNA vaccines expressing different forms of the SARS-CoV-2 S protein. The vaccine encoding the full-length S protein resulted in a dramatic reduction of viral replication in both the upper and lower respiratory tract after challenge with SARS-CoV-2. Van Doremalen et al. (118) found that vaccination of rhesus macaques with ChAdOx1 nCoV-19 induced balanced Th1/Th2 humoral and cellular immune responses, which prevent SARS-CoV-2 infection. Another vaccine developed by Gao et al. (119) is an inactivated SARS-CoV-2 virus vaccine (PiCoVacc), which induces the synthesis of SARS-CoV-2-specific neutralizing antibodies in macaques. In addition, the immunogenicity and protective efficacy of a single dose of adenovirus serotype 26 (Ad26) vector-based vaccines expressing the SARS-CoV-2 S protein were assessed in macaques. The single-shot Ad26 vaccine protected these animals against SARS-CoV-2 infection, and it is currently being evaluated in clinical trials (120).

African green monkeys are another NHP model for COVID-19 that replicates most facets of the human disease. These monkeys support high levels of SARS-CoV-2 replication and are susceptible to pronounced respiratory tract infection (121, 122). Importantly, subclinical infection of African green monkeys with SARS-CoV-2 resulted in prolonged shedding of infectious virus from both respiratory and gastrointestinal tracts (123).

A comparison of how human coronaviruses, including SARS-CoV, MERS-CoV, and SARS-CoV-2, develop in cynomolgus macaques showed that this NHP could be a predictive model to test preventive and therapeutic strategies against COVID-19 (124, 125). Salguero et al. (126) provided evidence that, though rhesus monkeys are considered to be the optimal model species, SARS-CoV-2 infection progresses similarly in both species. Guebre-Xabier et al. (127) revealed the efficacy of a SARS-CoV-2 subunit vaccine (NVX-CoV2373) in cynomolgus macaques. This

vaccine protected the animals against SARS-CoV-2 replication in both upper and lower respiratory tracts. In crab-eating macaques (*Macaca fascicularis*), serial computed tomography demonstrated that bilateral intrabronchial instillation of SARS-CoV-2 led to mild-to-moderate lung abnormalities (128).

Lu et al. (129) compared three species of NHPs, including rhesus macaques (*Macaca macaque*), crab-eating macaques (*Macaca fascicularis*), and common marmosets (*Callithrix jacchus*), that were challenged with the same strain of SARS-CoV-2 to identify the most suitable model for COVID-19. Based on analysis of clinical symptoms, viral replication, tissue tropism, and host responses to SARS-CoV-2 infection among the three species from two families of NHPs, rhesus macaques presented the strongest response to SARS-CoV-2. Hence, this animal closely recapitulated human-like conditions. Overall, NHPs are excellent animal models to better understand aspects related to human pathology in the current COVID-19 pandemic.

#### 4. COMPUTATIONAL AND MATHEMATICAL MODELS

Apart from the above-described in vivo and in vitro models, several computational and mathematical models are being used to study SARS-CoV-2 (130, 131). These computational models have been widely employed to screen potential drugs against SARS-CoV-2 (132, 133) and provide new components and targets for further in vitro and in vivo investigation of SARS-CoV-2 (134). Also, numerous computational and mathematical models predict the trend of COVID-19 transmission and number of infected individuals so as to help epidemic prevention and control measures (135–138).

#### 5. THERAPEUTIC APPROACHES FOR THE COVID-19 PANDEMIC

The global pandemic has created an urgent need to identify and validate therapeutic agents that can rapidly become part of clinical care. Antiviral drugs and host immunomodulators are the main approaches that may support recovery from COVID-19 (139). Antiviral agents can impair viral replication either by directly acting on viral targets or by targeting host proteins required for viral replication. Direct-acting antiviral compounds target SARS-CoV-2 proteins, including the S protein, proteases, helicase, and RNA-dependent RNA polymerase (RdRp) (140). Remdesivir (GS-5734), an inhibitor of RdRp with in vitro and in vivo inhibitory activity against SARS-CoV-2 (Table 5), was the first antiviral drug that received FDA approval in October 2020 for the treatment of COVID-19. This drug shortened the time to recovery in adults who were hospitalized with COVID-19 and showed evidence of lower respiratory tract infection (141). Later, in November 2020, WHO issued a conditional recommendation against the use of remdesivir in hospitalized patients, based on the lack of evidence that remdesivir improves survival and other outcomes in these patients (142).

Host-encoded proteins such as kinases and proteases employed by SARS-CoV-2 are potential targets of antiviral agents (143). Immunomodulatory agents, e.g., interleukin (IL)-1 (144) and IL-6 (145) receptor antagonists, might be beneficial for COVID-19 patients by attenuating an excessive host immune response. Based on evidence from several clinical trials (146), in September 2020, WHO issued a guideline on the use of dexamethasone and other corticosteroids (which have anti-inflammatory and immunosuppressant effects) as effective treatments for COVID-19. WHO recommends that corticosteroids (i.e., dexamethasone, hydrocortisone, or prednisone) be given orally or intravenously for the treatment of patients with severe and critical COVID-19.

The synthesis and investigation of potential new drugs is a time-consuming process that is not well suited for the COVID-19 pandemic. Accordingly, off-label use (repurposing) of existing

drugs is a promising approach, as it reduces development timelines and costs. Due to a lack of well-vetted guidelines, initial clinical use of such drugs in the COVID-19 pandemic has been uncoordinated. Some drugs were tested without a strong rationale or evidence to support their use against SARS-CoV-2. There remains a high demand to accelerate identification of agents with efficacy and safety for treating COVID-19. To this aim, both in vitro and in vivo experimental models of SARS-CoV-2 can elucidate aspects of pharmacology, toxicology, and immunology of the therapeutic strategies, which may evolve as understanding of COVID-19 biology improves (147). **Table 5** lists some repurposed antiviral drugs that have been tested in experimental models. It is clear that showing positive effects in a single nonhuman cell line assay cannot confirm the

**Table 5 Compounds, their mode of action, and treatment outcome in models of SARS-CoV-2**

Mechanism of action	Drug	Experimental model	Outcome	Reference(s)
Blockade of viral entry	Nafamostat	Vero E6	Inhibited SARS-CoV-2 infection	149
		Vero E6/TMPRSS2	Inhibited SARS-CoV-2 S protein-mediated fusion and SARS-CoV-2 infection in vitro in a cell type-dependent manner	150, 151
		Calu-3		
	Umifenovir (Arbidol)	Vero E6	Inhibited SARS-CoV-2 in vitro	152–154
Viral protease inhibitor	Lopinavir/ritonavir	Vero E6	Inhibited SARS-CoV-2 in vitro	152, 155–157
		Ferret	Marginally reduced overall clinical scores of infected ferrets but did not affect in vivo virus titers	106
Viral RNA polymerase inhibitor	Favipiravir	Vero E6	Not effective against SARS-CoV-2 infection in vitro	149, 155
			Rapid incorporation by the viral RNA polymerase complex resulted in SARS-CoV-2 lethal mutagenesis	158
		Syrian hamster	At high doses, antiviral activity in SARS-CoV-2-infected hamsters	159
	Remdesivir	Vero E6	Highly effective in the control of COVID-19 infection in vitro	149, 152, 155, 160
		Huh-7	Inhibited virus infection in human liver cancer cells	149
		Calu-3	Inhibited SARS-CoV-2 replication with submicromolar EC <sub>50</sub>	160
		Primary HAE	Inhibited SARS-CoV-2 replication	160
		Chimeric SARS-CoV encoding SARS-CoV-2 RNA polymerase in mice	Diminished lung viral load and improved pulmonary function compared to vehicle-treated animals	160
		Ad5-hACE2 transduced mice	Treatment 1 day prior to infection with continued dosing twice daily decreased weight loss, accelerated viral clearance, and diminished cellular infiltration in the lungs	78
		Rhesus macaques	Treatment initiated early during infection had clinical benefit in animals infected with SARS-CoV-2	114

(Continued)

**Table 5 (Continued)**

Mechanism of action	Drug	Experimental model	Outcome	Reference(s)
Miscellaneous mechanisms	Berberine	Vero E6	EC <sub>50</sub> in this model of SARS-CoV-2 infection was 10.6 $\mu$ M	152
	Chloroquine	Vero E6	Inhibited COVID-19	149, 152
		Calu-3	Did not inhibit infection of human lung cells with SARS-CoV-2	19
	Hydroxy-chloroquine	Vero	More potent than chloroquine in inhibiting SARS-CoV-2	161
		Vero E6	Inhibited SARS-CoV-2 infection	162
			Synergy with azithromycin on SARS-CoV-2	163
			Displayed antiviral activity	164
		Primary HAE	Lacked antiviral activity	164
		Ferret	Marginally reduced overall clinical scores but did not significantly affect in vivo viral titers	106
		Cynomolgus macaques	Data in this model do not support its use as an antiviral agent for the treatment of COVID-19 in humans	164
	Emetine	Vero E6	At 0.195 $\mu$ M in combination with remdesivir (6.25 $\mu$ M), there was 65% inhibition in viral load	155
			Off-target effects on acidophilic organelles (autophagosomes, lysosomes, endosomes) likely blunted SARS-CoV-2 replication	165

Abbreviations: COVID-19, coronavirus disease 2019; EC<sub>50</sub>, 50% maximal effective concentration; hACE2, human angiotensin-converting enzyme 2; HAE, human airway epithelial; SARS-CoV-2, severe acute respiratory syndrome coronavirus 2; TMPRSS2, transmembrane protease serine subtype 2.

efficacy of such drugs against COVID-19. A strong rationale and evidence (ideally in randomized controlled trials) are required for their use in COVID-19 patients.

Although several therapeutic approaches have been explored for COVID-19 treatment, to mitigate the influence of the virus on public health, society, and the economy, a vaccine is urgently needed to end the global COVID-19 pandemic. Currently, more than 200 vaccines are at various stages of development (148). These vaccine candidates comprise various platforms, including protein subunit, non-replicating or replicating viral vectors, DNA, inactivated virus, RNA, virus-like particle, and live attenuated virus. Many have entered clinical trials, and more than 10 vaccines are currently in Phase 3 or 2/3 clinical trials (**Table 6**). The FDA issued an emergency use authorization for the Pfizer-BioNTech and Moderna COVID-19 vaccines, and other effective and safe vaccines will likely be available soon.

## 6. CONCLUSION

This review summarizes models that are currently being used to study SARS-CoV-2 infection. We describe three main categories of in vitro, in vivo, and computational/mathematical models that each provide insights into different aspects of COVID-19. The in vitro models offer platforms to elucidate the viral life cycle, delineate disease pathology in tissue explants, and recapitulate host-virus interactions. Animal models include genetically modified mouse, Syrian hamster, ferret, and NHPs, with varying susceptibility to SARS-CoV-2 infection, which is likely largely dependent on species-specific differences in ACE2. These animal models are useful to investigate the effects of

**Table 6 Several vaccines that have been developed and used for COVID-19 treatment**

Vaccine	Vaccine Type	Developer	Animal model(s) tested	Clinical stage <sup>a</sup>	Reference(s)
BNT162	mRNA via lipid nanoparticles (lipid nanoparticle–formulated, nucleoside-modified mRNA vaccine that encodes the trimerized receptor-binding domain of the spike glycoprotein of SARS-CoV-2)	Pfizer/ BioNTech	Mice Rhesus macaques	Phase 2/3	166, 167, 195
mRNA-1273	mRNA via lipid nanoparticles (encoding the prefusion-stabilized spike protein of SARS-CoV-2)	Moderna	Rhesus macaques	Phase 3	168–170
PiCoVacc	Inactivated SARS-CoV-2 with aluminum hydroxide	Sinovac	BALB/c mice Wistar rats Rhesus macaques	Phase 3	119, 171
NVX-CoV2373	Full-length recombinant SARS-CoV-2 glycoprotein nanoparticle vaccine adjuvanted with Matrix M	Novavax	Cynomolgus macaques	Phase 3	127, 172
ChAdOx1 nCoV-19	Nonreplicating adenovirus (adenovirus vector–based vaccine, which encodes the spike protein of SARS-CoV-2)	AstraZeneca/ University of Oxford	Rhesus macaques	Phase 3	118, 173–175
Ad26COVS1	Nonreplicating adenovirus	Janssen Pharmaceuticals	BALB/c mice Hamsters Rhesus macaques	Phase 3	120, 176, 177
Adenovirus type 5 vector (Ad5CoV)	Nonreplicating adenovirus	CanSino Biological Inc./Beijing Institute of Biotechnology	NA	Phase 3	178, 179
rAd26-S + rAd5-S	Nonreplicating adenovirus	Gamaleya Research Institute	Nonhuman primates	Phase 3	180
BBV152 (Covaxin)	Inactivated SARS-CoV-2	Bharat Biotech International	Mice Rats Rabbits Hamsters Rhesus macaques	Phase 3	196–199
Inactivated vaccine	Inactivated SARS-CoV-2	Sinopharm/ Wuhan Institute of Biological Products	Rhesus macaques	Phase 3	181
Inactivated vaccine (BBIBP-CorV)	Inactivated SARS-CoV-2	Sinopharm/ Beijing Institute of Biological Products	Mice Rats Guinea pigs Rabbits Cynomolgus macaques Rhesus macaques	Phase 3	182, 183

<sup>a</sup>According to WHO Landscape as of December 22, 2020 (148).

Abbreviations: COVID-19, coronavirus disease 2019; mRNA, messenger RNA; NA, not applicable; SARS-CoV-2, severe acute respiratory syndrome coronavirus 2; WHO, World Health Organization.

a SARS-CoV-2 infection on specific target organs and to assess systemic effects of the infection. Computational and mathematical models are employed to predict COVID-19 transmission and to screen therapeutic strategies. By combining the outcomes of the in vitro, in vivo, and computational/mathematical models, one can identify fundamental aspects of COVID-19 in humans and

have useful platforms to test therapeutic agents and vaccines for safety and efficacy. These models can help elucidate aspects of pharmacology, toxicology, and immunology of repurposed antiviral drugs and aid in optimizing drug testing during the pandemic. No single animal model can fully recapitulate human pathogenesis and predict interventional responses in humans, but the models play a key role in assessing vaccines and therapeutics.

Even with the current clinical trials for vaccines, more work is urgently needed to understand host-infection interactions; to explain the extreme severity of the effects of infection, including perhaps the role of the nervous system in a neuroinflammatory response; and to develop effective treatments for those unable to be vaccinated or for whom vaccination only provides partial or no protection.

## DISCLOSURE STATEMENT

The authors are not aware of any affiliations, memberships, funding, or financial holdings that might be perceived as affecting the objectivity of this review.

## ACKNOWLEDGMENTS

This work was supported by Shiraz University of Medical Sciences (grant 22018). Partial support was also provided by Deutsche Forschungsgemeinschaft grant GRK 2174 and an International Brain Research Organization research fellowship award.

## LITERATURE CITED

1. Huang C, Wang Y, Li X, Ren L, Zhao J, et al. 2020. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* 395:497–506
2. Zhou P, Yang X-L, Wang X-G, Hu B, Zhang L, et al. 2020. A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature* 579:270–73
3. Hoffmann M, Kleine-Weber H, Schroeder S, Krüger N, Herrler T, et al. 2020. SARS-CoV-2 cell entry depends on ACE2 and TMPRSS2 and is blocked by a clinically proven protease inhibitor. *Cell* 181:271–80.e8
4. Leitner T, Kumar S. 2020. Where did SARS-CoV-2 come from? *Mol. Biol. Evol.* 37:2463–64
5. Salehi MS, Pandamooz S, Jurek B. 2020. Epidermal neural crest stem cells as a perspective for COVID-19 treatment. *Stem. Cell Rev. Rep.* <https://doi.org/10.1007/s12015-020-10028-3>
6. Sungnak W, Huang N, Becavin C, Berg M, Queen R, et al. 2020. SARS-CoV-2 entry factors are highly expressed in nasal epithelial cells together with innate immune genes. *Nat. Med.* 26:681–87
7. Pizzorno A, Padey B, Julien T, Trouillet-Assant S, Traversier A, et al. 2020. Characterization and treatment of SARS-CoV-2 in nasal and bronchial human airway epithelia. *Cell Rep. Med.* 1:100059
8. Sheahan TP, Sims AC, Zhou S, Graham RL, Pruijssers AJ, et al. 2020. An orally bioavailable broad-spectrum antiviral inhibits SARS-CoV-2 in human airway epithelial cell cultures and multiple coronaviruses in mice. *Sci. Transl. Med.* 12:eabb5883
9. Mulay A, Konda B, Garcia G Jr., Yao C, Beil S, et al. 2020. SARS-CoV-2 infection of primary human lung epithelium for COVID-19 modeling and drug discovery. *bioRxiv* 2020.06.29.174623. <https://doi.org/10.1101/2020.06.29.174623>
10. Hao S, Ning K, Kuz CA, Vorhies K, Yan Z, Qiu J. 2020. Long-term modeling of SARS-CoV-2 infection of in vitro cultured polarized human airway epithelium. *mBio* 11(6):e02852-20
11. Liu X, Wu Y, Rong L. 2020. Conditionally reprogrammed human normal airway epithelial cells at ALI: a physiological model for emerging viruses. *Virology* 535:280–89
12. Leist SR, Schäfer A, Martinez DR. 2020. Cell and animal models of SARS-CoV-2 pathogenesis and immunity. *Dis. Model. Mech.* 13:dmm046581

13. Matsuyama S, Nao N, Shirato K, Kawase M, Saito S, et al. 2020. Enhanced isolation of SARS-CoV-2 by TMPRSS2-expressing cells. *PNAS* 117:7001–3
14. Ogando NS, Dalebout TJ, Zevenhoven-Dobbe JC, Limpens R, van der Meer Y, et al. 2020. SARS-coronavirus-2 replication in Vero E6 cells: replication kinetics, rapid adaptation and cytopathology. *J. Gen. Virol.* 101:925–40
15. Harcourt J, Tamin A, Lu X, Kamili S, Sakthivel SK, et al. 2020. Severe acute respiratory syndrome coronavirus 2 from patient with coronavirus disease, United States. *Emerg. Infect. Dis.* 26:1266–73
16. Ellinger B, Bojkova D, Zaliani A, Cinatl J, Claussen C, et al. 2020. Identification of inhibitors of SARS-CoV-2 in-vitro cellular toxicity in human (Caco-2) cells using a large scale drug repurposing collection. *Res. Sq.* 23951. <https://www.researchsquare.com/article/rs-23951/v1>
17. Bojkova D, Klann K, Koch B, Widera M, Krause D, et al. 2020. Proteomics of SARS-CoV-2-infected host cells reveals therapy targets. *Nature* 583:469–72
18. Banerjee A, Nasir JA, Budyłowski P, Yip L, Aftanas P, et al. 2020. Isolation, sequence, infectivity, and replication kinetics of severe acute respiratory syndrome Coronavirus 2. *Emerg. Infect. Dis.* 26:2054–63
19. Hoffmann M, Mösbauer K, Hofmann-Winkler H, Kaul A, Kleine-Weber H, et al. 2020. Chloroquine does not inhibit infection of human lung cells with SARS-CoV-2. *Nature* 585:588–90
20. Ou X, Liu Y, Lei X, Li P, Mi D, et al. 2020. Characterization of spike glycoprotein of SARS-CoV-2 on virus entry and its immune cross-reactivity with SARS-CoV. *Nat. Commun.* 11:1620
21. Chu H, Chan JF-W, Yuen TT-T, Shuai H, Yuan S, et al. 2020. Comparative tropism, replication kinetics, and cell damage profiling of SARS-CoV-2 and SARS-CoV with implications for clinical manifestations, transmissibility, and laboratory studies of COVID-19: an observational study. *Lancet Microbe* 1:e14–23
22. Cagno V. 2020. SARS-CoV-2 cellular tropism. *Lancet Microbe* 1:e2–3
23. Ma-Lauer Y, Carbajo-Lozoya J, Hein MY, Müller MA, Deng W, et al. 2016. p53 down-regulates SARS coronavirus replication and is targeted by the SARS-unique domain and PLpro via E3 ubiquitin ligase RCHY1. *PNAS* 113:E5192–201
24. Yang L, Han Y, Nilsson-Payant BE, Gupta V, Wang P, et al. 2020. A human pluripotent stem cell-based platform to study SARS-CoV-2 tropism and model virus infection in human cells and organoids. *Cell Stem Cell* 27:125–36.e7
25. Surendran H, Nandakumar S, Pal R. 2020. Human induced pluripotent stem cell-derived lung epithelial system for SARS-CoV-2 infection modeling and its potential in drug repurposing. *Stem Cells Dev.* 29(21):1365–69
26. Huang J, Hume AJ, Abo KM, Werder RB, Villacorta-Martin C, et al. 2020. SARS-CoV-2 infection of pluripotent stem cell-derived human lung alveolar type 2 cells elicits a rapid epithelial-intrinsic inflammatory response. *Cell Stem Cell* 27:962–73.e7
27. Abo KM, Ma L, Matte T, Huang J, Alysandratos KD, et al. 2020. Human iPSC-derived alveolar and airway epithelial cells can be cultured at air-liquid interface and express SARS-CoV-2 host factors. *bioRxiv* 2020.06.03.132639. <https://doi.org/10.1101/2020.06.03.132639>
28. Duan F, Guo L, Yang L, Han Y, Thakur A, et al. 2020. Modeling COVID-19 with human pluripotent stem cell-derived cells reveals synergistic effects of anti-inflammatory macrophages with ACE2 inhibition against SARS-CoV-2. *Res. Sq.* 62758. <https://www.researchsquare.com/article/rs-62758/v1>
29. Bojkova D, Wagner JUG, Shumliakivska M, Aslan GS, Saleem U, et al. 2020. SARS-CoV-2 infects and induces cytotoxic effects in human cardiomyocytes. *Cardiovasc. Res.* 116(14):2207–15
30. Sharma A, Garcia G Jr., Wang Y, Plummer JT, Morizono K, et al. 2020. Human iPSC-derived cardiomyocytes are susceptible to SARS-CoV-2 infection. *Cell Rep. Med.* 1:100052
31. Pérez-Bermejo JA, Kang S, Rockwood SJ, Simoneau CR, Joy DA, et al. 2020. SARS-CoV-2 infection of human iPSC-derived cardiac cells predicts novel cytopathic features in hearts of COVID-19 patients. *bioRxiv* 2020.08.25.265561. <https://www.biorxiv.org/content/10.1101/2020.08.25.265561v2>
32. Mao L, Jin H, Wang M, Hu Y, Chen S, et al. 2020. Neurologic manifestations of hospitalized patients with coronavirus disease 2019 in Wuhan, China. *JAMA Neurol.* 77:683–90
33. Mao X-Y, Jin W-L. 2020. iPSCs-derived platform: a feasible tool for probing the neurotropism of SARS-CoV-2. *ACS Chem. Neurosci.* 11:2489–91
34. Zhang B-Z, Chu H, Han S, Shuai H, Deng J, et al. 2020. SARS-CoV-2 infects human neural progenitor cells and brain organoids. *Cell Res.* 30:928–31

35. Jacob F, Pather SR, Huang W-K, Zhang F, Wong SZH, et al. 2020. Human pluripotent stem cell-derived neural cells and brain organoids reveal SARS-CoV-2 neurotropism predominates in choroid plexus epithelium. *Cell Stem Cell*. 27(6):937–50.e9
36. Kim J, Koo B-K, Knoblich JA. 2020. Human organoids: model systems for human biology and medicine. *Nat. Rev. Mol. Cell Biol.* 21:571–84
37. Clevers H. 2020. COVID-19: organoids go viral. *Nat. Rev. Mol. Cell Biol.* 21:355–56
38. Mallapaty S. 2020. Mini organs reveal how the coronavirus ravages the body. *Nature* 583:15–16
39. Elbadawi M, Efferth T. 2020. Organoids of human airways to study infectivity and cytopathy of SARS-CoV-2. *Lancet Respir. Med.* 8:E55–56
40. Bose B. 2020. Induced pluripotent stem cells (iPSCs) derived 3D human lung organoids from different ethnicities to understand the SARS-CoV2 severity/infectivity percentage. *Stem. Cell Rev. Rep.* 17:293–95
41. Han Y, Duan X, Yang L, Nilsson-Payant BE, Wang P, et al. 2020. Identification of SARS-CoV-2 inhibitors using lung and colonic organoids. *Nature* 589:270–75
42. Suzuki T, Itoh Y, Sakai Y, Saito A, Okuzaki D, et al. 2020. Generation of human bronchial organoids for SARS-CoV-2 research. bioRxiv 2020.05.25.115600. <https://www.biorxiv.org/content/10.1101/2020.05.25.115600v2>
43. Gu H, Yuan G. 2020. Identification of potential key genes for SARS-CoV-2 infected human bronchial organoids based on bioinformatics analysis. bioRxiv 2020.08.18.256735. <https://www.biorxiv.org/content/10.1101/2020.08.18.256735v2>
44. Kim HK, Kim H, Lee MK, Choi WH, Jang Y, et al. 2020. Generation of tonsil organoids as an ex vivo model for SARS-CoV-2 infection. bioRxiv 2020.08.06.239574. <https://www.biorxiv.org/content/10.1101/2020.08.06.239574v1>
45. Dickson I. 2020. Organoids demonstrate gut infection by SARS-CoV-2. *Nat. Rev. Gastroenterol. Hepatol.* 17:383
46. Zhang H, Kang Z, Gong H, Xu D, Wang J, et al. 2020. Digestive system is a potential route of COVID-19: an analysis of single-cell coexpression pattern of key proteins in viral entry process. *Gut* 69:1010–18
47. Lamers MM, Beumer J, van der Vaart J, Knoops K, Puschhof J, et al. 2020. SARS-CoV-2 productively infects human gut enterocytes. *Science* 369:50–54
48. Zhou J, Li C, Liu X, Chiu MC, Zhao X, et al. 2020. Infection of bat and human intestinal organoids by SARS-CoV-2. *Nat. Med.* 26:1077–83
49. Stanifer ML, Kee C, Cortese M, Zumaran CM, Triana S, et al. 2020. Critical role of type III interferon in controlling SARS-CoV-2 infection in human intestinal epithelial cells. *Cell Rep.* 32:107863
50. Xu L, Liu J, Lu M, Yang D, Zheng X. 2020. Liver injury during highly pathogenic human coronavirus infections. *Liver Int.* 40:998–1004
51. Fan Z, Chen L, Li J, Cheng X, Yang J, et al. 2020. Clinical features of COVID-19-related liver functional abnormality. *Clin. Gastroenterol. Hepatol.* 18:1561–66
52. Qi F, Qian S, Zhang S, Zhang Z. 2020. Single cell RNA sequencing of 13 human tissues identify cell types and receptors of human coronaviruses. *Biochem. Biophys. Res. Commun.* 526:135–40
53. Chai X, Hu L, Zhang Y, Han W, Lu Z, et al. 2020. Specific ACE2 expression in cholangiocytes may cause liver damage after 2019-nCoV infection. bioRxiv 2020.02.03.931766. <https://www.biorxiv.org/content/10.1101/2020.02.03.931766v1>
54. Zhao B, Ni C, Gao R, Wang Y, Yang L, et al. 2020. Recapitulation of SARS-CoV-2 infection and cholangiocyte damage with human liver ductal organoids. *Protein Cell* 11:771–75
55. Monteil V, Kwon H, Prado P, Hagelkrüys A, Wimmer RA, et al. 2020. Inhibition of SARS-CoV-2 infections in engineered human tissues using clinical-grade soluble human ACE2. *Cell* 181:905–13.e7
56. Iadecola C, Anrather J, Kamel H. 2020. Effects of COVID-19 on the nervous system. *Cell* 183:16–27
57. Bullen CK, Hogberg HT, Bahadiri-Talbot A, Bishai WR, Hartung T, et al. 2020. Infectability of human BrainSphere neurons suggests neurotropism of SARS-CoV-2. *Altex* 37:665–71
58. Yi SA, Nam KH, Yun J, Gim D, Joe D, et al. 2020. Infection of brain organoids and 2D cortical neurons with SARS-CoV-2 pseudovirus. *Viruses* 12:1004
59. Song E, Zhang C, Israelow B, Lu-Culligan A, Vieites Prado A, et al. 2020. Neuroinvasion of SARS-CoV-2 in human and mouse brain. bioRxiv 2020.06.25.169946. <https://www.biorxiv.org/content/10.1101/2020.06.25.169946v2>

60. Ramani A, Müller L, Ostermann PN, Gabriel E, Abida-Islam P, et al. 2020. SARS-CoV-2 targets neurons of 3D human brain organoids. *EMBO J.* 39:e106230
61. Tang H, Abouleila Y, Si L, Ortega-Prieto AM, Mummery CL, et al. 2020. Human organs-on-chips for virology. *Trends Microbiol.* 28:934–46
62. Si L, Bai H, Rodas M, Cao W, Oh CY, et al. 2020. Human organ chip-enabled pipeline to rapidly repurpose therapeutics during viral pandemics. *bioRxiv* 2020.04.13.039917. <https://www.biorxiv.org/content/10.1101/2020.04.13.039917v3>
63. Thacker V, Sharma K, Dhar N, Mancini G-F, Sordet-Dessimoz J, McKinney JD. 2020. Rapid endothelial infection, endothelialitis and vascular damage characterise SARS-CoV-2 infection in a human lung-on-chip model. *bioRxiv* 2020.08.10.243220. <https://www.biorxiv.org/content/10.1101/2020.08.10.243220v2>
64. Zhang M, Wang P, Luo R, Wang Y, Li Z, et al. 2020. Biomimetic human disease model of SARS-CoV-2 induced lung injury and immune responses on organ chip system. *Adv. Sci.* 8:2002928
65. Guo Y, Luo R, Wang Y, Deng P, Zhang M, et al. 2020. Modeling SARS-CoV-2 infection in vitro with a human intestine-on-chip device. *bioRxiv* 2020.09.01.277780. <https://www.biorxiv.org/content/10.1101/2020.09.01.277780v1>
66. Grivel J-C, Margolis L. 2009. Use of human tissue explants to study human infectious agents. *Nat. Protoc.* 4:256–69
67. Pandamooz S, Nabiuni M, Miyan J, Ahmadiani A, Dargahi L. 2016. Organotypic spinal cord culture: a proper platform for the functional screening. *Mol. Neurobiol.* 53:4659–74
68. Chu H, Chan JF-W, Wang Y, Yuen TT-T, Chai Y, et al. 2020. Comparative replication and immune activation profiles of SARS-CoV-2 and SARS-CoV in human lungs: an ex vivo study with implications for the pathogenesis of COVID-19. *Clin. Infect. Dis.* 71:1400–9
69. Ravindra NG, Alfajaro MM, Gasque V, Habet V, Wei J, et al. 2020. Single-cell longitudinal analysis of SARS-CoV-2 infection in human airway epithelium. *bioRxiv* 2020.05.06.081695. <https://www.biorxiv.org/content/10.1101/2020.05.06.081695v2>
70. Hui KPY, Cheung M-C, Perera RAPM, Ng K-C, Bui CHT, et al. 2020. Tropism, replication competence, and innate immune responses of the coronavirus SARS-CoV-2 in human respiratory tract and conjunctiva: an analysis in ex-vivo and in-vitro cultures. *Lancet Respir. Med.* 8:687–95
71. Zhu N, Wang W, Liu Z, Liang C, Wang W, et al. 2020. Morphogenesis and cytopathic effect of SARS-CoV-2 infection in human airway epithelial cells. *Nat. Commun.* 11:3910
72. Johansen M, Irving A, Montagutelli X, Tate M, Rudloff I, et al. 2020. Animal and translational models of SARS-CoV-2 infection and COVID-19. *Mucosal Immunol.* 13:877–91
73. Singh A, Singh RS, Sarma P, Batra G, Joshi R, et al. 2020. A comprehensive review of animal models for coronaviruses: SARS-CoV-2, SARS-CoV, and MERS-CoV. *Virol. Sin.* 35:290–304
74. Neerukonda SN, Katneni U. 2020. A review on SARS-CoV-2 virology, pathophysiology, animal models, and anti-viral interventions. *Pathogens* 9:426
75. Takayama K. 2020. In vitro and animal models for SARS-CoV-2 research. *Trends Pharmacol. Sci.* 41:513–17
76. Letko M, Marzi A, Munster V. 2020. Functional assessment of cell entry and receptor usage for SARS-CoV-2 and other lineage B betacoronaviruses. *Nat. Microbiol.* 5:562–69
77. Wan Y, Shang J, Graham R, Baric RS, Li F. 2020. Receptor recognition by the novel coronavirus from Wuhan: an analysis based on decade-long structural studies of SARS coronavirus. *J. Virol.* 94:e00127–20
78. Sun J, Zhuang Z, Zheng J, Li K, Wong RL-Y, et al. 2020. Generation of a broadly useful model for COVID-19 pathogenesis, vaccination, and treatment. *Cell* 182:734–43.e5
79. Bao L, Deng W, Huang B, Gao H, Liu J, et al. 2020. The pathogenicity of SARS-CoV-2 in hACE2 transgenic mice. *Nature* 583:830–33
80. Winkler ES, Bailey AL, Kafai NM, Nair S, McCune BT, et al. 2020. SARS-CoV-2 infection of human ACE2-transgenic mice causes severe lung inflammation and impaired function. *Nat. Immunol.* 21:1327–35
81. Jiang R-D, Liu M-Q, Chen Y, Shan C, Zhou Y-W, et al. 2020. Pathogenesis of SARS-CoV-2 in transgenic mice expressing human angiotensin-converting enzyme 2. *Cell* 182:50–58

82. Sun S-H, Chen Q, Gu H-J, Yang G, Wang Y-X, et al. 2020. A mouse model of SARS-CoV-2 infection and pathogenesis. *Cell Host Microbe* 28:124–33.e4
83. Hassan AO, Case JB, Winkler ES, Thackray LB, Kafai NM, et al. 2020. A SARS-CoV-2 infection model in mice demonstrates protection by neutralizing antibodies. *Cell* 182:744–53.e4
84. Pujhari S, Rasgon JL. 2020. Mice with humanized-lungs and immune system—an idealized model for COVID-19 and other respiratory illness. *Virulence* 11:486–88
85. Wahl A, De C, Fernandez MA, Lenarcic EM, Xu Y, et al. 2019. Precision mouse models with expanded tropism for human pathogens. *Nat. Biotechnol.* 37:1163–73
86. Mori M, Furuhashi K, Danielsson JA, Hirata Y, Kakiuchi M, et al. 2019. Generation of functional lungs via conditional blastocyst complementation using pluripotent stem cells. *Nat. Med.* 25:1691–98
87. Gu H, Chen Q, Yang G, He L, Fan H, et al. 2020. Adaptation of SARS-CoV-2 in BALB/c mice for testing vaccine efficacy. *Science* 369:1603–7
88. Dinno KH, Leist SR, Schäfer A, Edwards CE, Martinez DR, et al. 2020. A mouse-adapted model of SARS-CoV-2 to test COVID-19 countermeasures. *Nature* 586:560–66
89. Lau SY, Wang P, Mok BW, Zhang AJ, Chu H, et al. 2020. Attenuated SARS-CoV-2 variants with deletions at the S1/S2 junction. *Emerg. Microbes Infect.* 9:837–42
90. Chan JF-W, Yuan S, Zhang AJ, Poon VK-M, Chan CC-S, et al. 2020. Surgical mask partition reduces the risk of non-contact transmission in a golden Syrian hamster model for Coronavirus Disease 2019 (COVID-19). *Clin. Infect. Dis.* 71:2139–49
91. Schaecher SR, Stabenow J, Oberle C, Schriewer J, Buller RM, et al. 2008. An immunosuppressed Syrian golden hamster model for SARS-CoV infection. *Virology* 380:312–21
92. Chan JF-W, Zhang AJ, Yuan S, Poon VK-M, Chan CC-S, et al. 2020. Simulation of the clinical and pathological manifestations of Coronavirus Disease 2019 (COVID-19) in golden Syrian hamster model: implications for disease pathogenesis and transmissibility. *Clin. Infect. Dis.* 71(9):2428–446
93. Imai M, Iwatsuki-Horimoto K, Hatta M, Loeber S, Halfmann PJ, et al. 2020. Syrian hamsters as a small animal model for SARS-CoV-2 infection and countermeasure development. *PNAS* 117:16587–95
94. Sia SF, Yan L-M, Chin AW, Fung K, Choy K-T, et al. 2020. Pathogenesis and transmission of SARS-CoV-2 in golden hamsters. *Nature* 583:834–38
95. Rogers TF, Zhao F, Huang D, Beutler N, Burns A, et al. 2020. Isolation of potent SARS-CoV-2 neutralizing antibodies and protection from disease in a small animal model. *Science* 369:956–63
96. Subbarao K. 2020. SARS-CoV-2: A new song recalls an old melody. *Cell Host Microbe* 27:692–94
97. Osterrieder N, Bertzbach LD, Dietert K, Abdelgawad A, Vladimirova D, et al. 2020. Age-dependent progression of SARS-CoV-2 infection in Syrian hamsters. *Viruses* 12:779
98. Boudewijns R, Thibaut HJ, Kaptein SJF, Li R, Vergote V, et al. 2020. STAT2 signaling restricts viral dissemination but drives severe pneumonia in SARS-CoV-2 infected hamsters. *Nat. Commun.* 11:5838
99. Suresh V, Parida D, Minz AP, Senapati S. 2020. Tissue distribution of ACE2 protein in Syrian golden hamster (*Mesocricetus auratus*) and its possible implications in SARS-CoV-2 related studies. *bioRxiv* 2020.06.29.177154. <https://www.biorxiv.org/content/10.1101/2020.06.29.177154v1>
100. Martina BE, Haagmans BL, Kuiken T, Fouchier RA, Rimmelzwaan GF, et al. 2003. SARS virus infection of cats and ferrets. *Nature* 425:915
101. Chu Y-K, Ali GD, Jia F, Li Q, Kelvin D, et al. 2008. The SARS-CoV ferret model in an infection–challenge study. *Virology* 374:151–63
102. Shi J, Wen Z, Zhong G, Yang H, Wang C, et al. 2020. Susceptibility of ferrets, cats, dogs, and other domesticated animals to SARS-coronavirus 2. *Science* 368:1016–20
103. Schlottau K, Rissmann M, Graaf A, Schön J, Sehl J, et al. 2020. SARS-CoV-2 in fruit bats, ferrets, pigs, and chickens: an experimental transmission study. *Lancet Microbe* 1:e218–25
104. Kim Y-I, Kim S-G, Kim S-M, Kim E-H, Park S-J, et al. 2020. Infection and rapid transmission of SARS-CoV-2 in ferrets. *Cell Host Microbe* 27:704–9.e2
105. Richard M, Kok A, de Meulder D, Bestebroer TM, Lamers MM, et al. 2020. SARS-CoV-2 is transmitted via contact and via the air between ferrets. *Nat. Commun.* 11:3496
106. Park S-J, Yu K-M, Kim Y-I, Kim S-M, Kim E-H, et al. 2020. Antiviral efficacies of FDA-approved drugs against SARS-CoV-2 infection in ferrets. *mBio* 11:e01114-20

107. Liu H-L, Yeh IJ, Phan NN, Wu Y-H, Yen M-C, et al. 2020. Gene signatures of SARS-CoV/SARS-CoV-2-infected ferret lungs in short- and long-term models. *Infect. Genet. Evol.* 85:104438
108. Santos WJ, Guiraldi LM, Lucheis SB. 2020. Should we be concerned about COVID-19 with nonhuman primates? *Am. J. Primatol.* 82:e23158
109. Rowe T, Gao G, Hogan RJ, Crystal RG, Voss TG, et al. 2004. Macaque model for severe acute respiratory syndrome. *J. Virol.* 78:11401–4
110. Melin AD, Janiak MC, Marrone F, Arora PS, Higham JP. 2020. Comparative ACE2 variation and primate COVID-19 risk. *Commun. Biol.* 3:641
111. Zhao X, Chen D, Szabla R, Zheng M, Li G, et al. 2020. Broad and differential animal angiotensin-converting enzyme 2 receptor usage by SARS-CoV-2. *J. Virol.* 94:e00940-20
112. Shan C, Yao Y-F, Yang X-L, Zhou Y-W, Gao G, et al. 2020. Infection with novel coronavirus (SARS-CoV-2) causes pneumonia in Rhesus macaques. *Cell Res.* 30:670–77
113. Munster VJ, Feldmann F, Williamson BN, van Doremalen N, Pérez-Pérez L, et al. 2020. Respiratory disease in rhesus macaques inoculated with SARS-CoV-2. *Nature* 585:268–72
114. Williamson BN, Feldmann F, Schwarz B, Meade-White K, Porter DP, et al. 2020. Clinical benefit of remdesivir in rhesus macaques infected with SARS-CoV-2. *Nature* 585:273–76
115. Deng W, Bao L, Liu J, Xiao C, Liu J, et al. 2020. Primary exposure to SARS-CoV-2 protects against reinfection in rhesus macaques. *Science* 369:818–23
116. Chandrashekar A, Liu J, Martinot AJ, McMahan K, Mercado NB, et al. 2020. SARS-CoV-2 infection protects against rechallenge in rhesus macaques. *Science* 369:812–17
117. Yu J, Tostanoski LH, Peter L, Mercado NB, McMahan K, et al. 2020. DNA vaccine protection against SARS-CoV-2 in rhesus macaques. *Science* 369:806–11
118. van Doremalen N, Lambe T, Spencer A, Belij-Rammerstorfer S, Purushotham JN, et al. 2020. ChAdOx1 nCoV-19 vaccine prevents SARS-CoV-2 pneumonia in rhesus macaques. *Nature* 586:578–82
119. Gao Q, Bao L, Mao H, Wang L, Xu K, et al. 2020. Development of an inactivated vaccine candidate for SARS-CoV-2. *Science* 369:77–81
120. Mercado NB, Zahn R, Wegmann F, Loos C, Chandrashekar A, et al. 2020. Single-shot Ad26 vaccine protects against SARS-CoV-2 in rhesus macaques. *Nature* 586:583–88
121. Cross RW, Agans KN, Prasad AN, Borisevich V, Woolsey C, et al. 2020. Intranasal exposure of African green monkeys to SARS-CoV-2 results in acute phase pneumonia with shedding and lung injury still present in the early convalescence phase. *J. Virol.* 7. 17:125
122. Woolsey C, Borisevich V, Prasad AN, Agans KN, Deer DJ, et al. 2021. Establishment of an African green monkey model for COVID-19 and protection against re-infection. *Nat. Immunol.* 22:86–98
123. Hartman AL, Nambulli S, McMillen CM, White AG, Tilston-Lunel NL, et al. 2020. SARS-CoV-2 infection of African green monkeys results in mild respiratory disease discernible by PET/CT imaging and shedding of infectious virus from both respiratory and gastrointestinal tracts. *PLOS Pathog.* 16:e1008903
124. Rockx B, Kuiken T, Herfst S, Bestebroer T, Lamers MM, et al. 2020. Comparative pathogenesis of COVID-19, MERS, and SARS in a nonhuman primate model. *Science* 368:1012–15
125. Le Bras A. 2020. SARS-CoV-2 causes COVID-19-like disease in cynomolgus macaques. *Lab Anim.* 49:174
126. Salguero FJ, White AD, Slack GS, Fotheringham SA, Bewley KR, et al. 2020. Comparison of Rhesus and Cynomolgus macaques as an authentic model for COVID-19. *bioRxiv* 2020.09.17.301093. <https://www.biorxiv.org/content/10.1101/2020.09.17.301093v1>
127. Guebre-Xabier M, Patel N, Tian J-H, Zhou B, Maciejewski S, et al. 2020. NVX-CoV2373 vaccine protects cynomolgus macaque upper and lower airways against SARS-CoV-2 challenge. *Vaccine* 38:7892–96
128. Finch CL, Crozier I, Lee JH, Byrum R, Cooper TK, et al. 2020. Characteristic and quantifiable COVID-19-like abnormalities in CT- and PET/CT-imaged lungs of SARS-CoV-2-infected crab-eating macaques (*Macaca fascicularis*). *bioRxiv* 2020.05.14.096727. <https://www.biorxiv.org/content/10.1101/2020.05.14.096727v1>
129. Lu S, Zhao Y, Yu W, Yang Y, Gao J, et al. 2020. Comparison of nonhuman primates identified the suitable model for COVID-19. *Signal. Transduct. Target. Ther.* 5:157

130. Machado JAT, Rocha-Neves JM, Andrade JP. 2020. Computational analysis of the SARS-CoV-2 and other viruses based on the Kolmogorov's complexity and Shannon's information theories. *Nonlinear Dyn.* 101:1731–50
131. Ostaszewski M, Mazein A, Gillespie ME, Kuperstein I, Niarakis A, et al. 2020. COVID-19 disease map, building a computational repository of SARS-CoV-2 virus-host interaction mechanisms. *Sci. Data* 7:136
132. Ghosh K, Amin SA, Gayen S, Jha T. 2021. Chemical-informatics approach to COVID-19 drug discovery: exploration of important fragments and data mining based prediction of some hits from natural origins as main protease (Mpro) inhibitors. *J. Mol. Struct.* 1224:129026
133. Bobrowski T, Alves V, Melo-Filho CC, Korn D, Auerbach SS, et al. 2020. Computational models identify several FDA approved or experimental drugs as putative agents against SARS-CoV-2. ChemRxiv 12153594. <https://doi.org/10.26434/chemrxiv.12153594.v1>
134. Wu C, Liu Y, Yang Y, Zhang P, Zhong W, et al. 2020. Analysis of therapeutic targets for SARS-CoV-2 and discovery of potential drugs by computational methods. *Acta Pharm. Sin. B* 10:766–88
135. Serhani M, Labbardi H. 2021. Mathematical modeling of COVID-19 spreading with asymptomatic infected and interacting peoples. *J. Appl. Math. Comput.* In press. <https://doi.org/10.1007/s12190-020-01421-9>
136. Torrealba-Rodriguez O, Conde-Gutierrez RA, Hernandez-Javier AL. 2020. Modeling and prediction of COVID-19 in Mexico applying mathematical and computational models. *Chaos Solitons Fractals* 138:109946
137. Ndairou F, Area I, Nieto JJ, Torres DFM. 2020. Mathematical modeling of COVID-19 transmission dynamics with a case study of Wuhan. *Chaos Solitons Fractals* 135:109846
138. Jiang S, Li Q, Li C, Liu S, He X, et al. 2020. Mathematical models for devising the optimal SARS-CoV-2 strategy for eradication in China, South Korea, and Italy. *J. Transl. Med.* 18:345
139. Simonis A, Theobald SJ, Fatkenheuer G, Rybníček J, Malin JJ. 2020. A comparative analysis of remdesivir and other repurposed antivirals against SARS-CoV-2. *EMBO Mol. Med.* 13:e13105
140. Grobler JA, Anderson AS, Fernandes P, Diamond MS, Colvis CM, et al. 2020. Accelerated preclinical paths to support rapid development of COVID-19 therapeutics. *Cell Host Microbe* 28:638–45
141. Beigel JH, Tomashek KM, Dodd LE, Mehta AK, Zingman BS, et al. 2020. Remdesivir for the treatment of Covid-19—final report. *N. Engl. J. Med.* 383:1813–26
142. WHO Solidarity Trial Consort. 2020. Repurposed antiviral drugs for Covid-19—interim WHO solidarity trial results. *N. Engl. J. Med.* 2020:NEJMoa2023184
143. Bouhaddou M, Memon D, Meyer B, White KM, Rezelj VV, et al. 2020. The global phosphorylation landscape of SARS-CoV-2 infection. *Cell* 182:685–712.e19
144. Cauchois R, Koubi M, Delarbre D, Manet C, Carvelli J, et al. 2020. Early IL-1 receptor blockade in severe inflammatory respiratory failure complicating COVID-19. *PNAS* 117:18951–53
145. Guillen L, Padilla S, Fernandez M, Agullo V, Garcia JA, et al. 2020. Preemptive interleukin-6 blockade in patients with COVID-19. *Sci. Rep.* 10:16826
146. RECOVERY Collab. Group. 2020. Dexamethasone in hospitalized patients with Covid-19—preliminary report. *N. Engl. J. Med.* 2020:NEJMoa2021436
147. Hewitt JA, Lutz C, Florence WC, Pitt MLM, Rao S, et al. 2020. ACTIVating resources for the COVID-19 pandemic: in vivo models for vaccines and therapeutics. *Cell Host Microbe* 28:646–59
148. WHO (World Health Organ.). 2020. *DRAFT landscape of COVID-19 candidate vaccines*. Database, WHO, Geneva. <https://www.who.int/publications/m/item/draft-landscape-of-covid-19-candidate-vaccines>
149. Wang M, Cao R, Zhang L, Yang X, Liu J, et al. 2020. Remdesivir and chloroquine effectively inhibit the recently emerged novel coronavirus (2019-nCoV) in vitro. *Cell Res.* 30:269–71
150. Hoffmann M, Schroeder S, Kleine-Weber H, Müller MA, Drosten C, Pohlmann S. 2020. Nafamostat mesylate blocks activation of SARS-CoV-2: new treatment option for COVID-19. *Antimicrob. Agents Chemother.* 64(6):e00754-20
151. Yamamoto M, Kiso M, Sakai-Tagawa Y, Iwatsuki-Horimoto K, Imai M, et al. 2020. The anticoagulant nafamostat potently inhibits SARS-CoV-2 S protein-mediated fusion in a cell fusion assay system and viral infection in vitro in a cell-type-dependent manner. *Viruses* 12:629

152. Pizzorno A, Padey B, Dubois J, Julien T, Traversier A, et al. 2020. In vitro evaluation of antiviral activity of single and combined repurposable drugs against SARS-CoV-2. *Antiviral Res.* 181:104878
153. Touret F, Gilles M, Barral K, Nougairede A, van Helden J, et al. 2020. In vitro screening of a FDA approved chemical library reveals potential inhibitors of SARS-CoV-2 replication. *Sci. Rep.* 10:13093
154. Wang X, Cao R, Zhang H, Liu J, Xu M, et al. 2020. The anti-influenza virus drug, arbidol is an efficient inhibitor of SARS-CoV-2 in vitro. *Cell Discov.* 6:28
155. Choy KT, Wong AY, Kaewpreedee P, Sia SF, Chen D, et al. 2020. Remdesivir, lopinavir, emetine, and homoharringtonine inhibit SARS-CoV-2 replication in vitro. *Antiviral Res.* 178:104786
156. Kang CK, Seong MW, Choi SJ, Kim TS, Choe PG, et al. 2020. In vitro activity of lopinavir/ritonavir and hydroxychloroquine against severe acute respiratory syndrome coronavirus 2 at concentrations achievable by usual doses. *Korean J. Intern. Med.* 35:782–87
157. Zhang L, Liu J, Cao R, Xu M, Wu Y, et al. 2020. Comparative antiviral efficacy of viral protease inhibitors against the novel SARS-CoV-2 in vitro. *Virol. Sin.* 35:776–84
158. Shannon A, Selisko B, Le NT, Huchting J, Touret F, et al. 2020. Rapid incorporation of favipiravir by the fast and permissive viral RNA polymerase complex results in SARS-CoV-2 lethal mutagenesis. *Nat. Commun.* 11:4682
159. Kaptein SJF, Jacobs S, Langendries L, Seldeslachts L, Ter Horst S, et al. 2020. Favipiravir at high doses has potent antiviral activity in SARS-CoV-2-infected hamsters, whereas hydroxychloroquine lacks activity. *PNAS* 117:26955–65
160. Pruijssers AJ, George AS, Schafer A, Leist SR, Gralinski LE, et al. 2020. Remdesivir inhibits SARS-CoV-2 in human lung cells and chimeric SARS-CoV expressing the SARS-CoV-2 RNA polymerase in mice. *Cell Rep.* 32:107940
161. Yao X, Ye F, Zhang M, Cui C, Huang B, et al. 2020. In vitro antiviral activity and projection of optimized dosing design of hydroxychloroquine for the treatment of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). *Clin. Infect. Dis.* 71:732–39
162. Liu J, Cao R, Xu M, Wang X, Zhang H, et al. 2020. Hydroxychloroquine, a less toxic derivative of chloroquine, is effective in inhibiting SARS-CoV-2 infection in vitro. *Cell Discov.* 6:16
163. Andreani J, Le Bideau M, Duflot I, Jardot P, Rolland C, et al. 2020. In vitro testing of combined hydroxychloroquine and azithromycin on SARS-CoV-2 shows synergistic effect. *Microb. Pathog.* 145:104228
164. Maisonnasse P, Guedj J, Contreras V, Behillil S, Solas C, et al. 2020. Hydroxychloroquine use against SARS-CoV-2 infection in non-human primates. *Nature* 585:584–87
165. Sauvat A, Ciccosanti F, Colavita F, Di Rienzo M, Castilletti C, et al. 2020. On-target versus off-target effects of drugs inhibiting the replication of SARS-CoV-2. *Cell Death Dis.* 11:656
166. Sahin U, Muik A, Derhovanessian E, Vogler I, Kranz LM, et al. 2020. COVID-19 vaccine BNT162b1 elicits human antibody and TH1 T cell responses. *Nature* 586:594–99
167. Mulligan MJ, Lyke KE, Kitchin N, Absalon J, Gurtman A, et al. 2020. Phase I/II study of COVID-19 RNA vaccine BNT162b1 in adults. *Nature* 586:589–93
168. Corbett KS, Flynn B, Foulds KE, Francica JR, Boyoglu-Barnum S, et al. 2020. Evaluation of the mRNA-1273 vaccine against SARS-CoV-2 in nonhuman primates. *N. Engl. J. Med.* 383:1544–55
169. Jackson LA, Anderson EJ, Roupheal NG, Roberts PC, Makhene M, et al. 2020. An mRNA vaccine against SARS-CoV-2—preliminary report. *N. Engl. J. Med.* 383:1920–31
170. Anderson EJ, Roupheal NG, Widge AT, Jackson LA, Roberts PC, et al. 2020. Safety and immunogenicity of SARS-CoV-2 mRNA-1273 vaccine in older adults. *N. Engl. J. Med.* 383:2427–38
171. Zhang Y, Zeng G, Pan H, Li C, Hu Y, et al. 2020. Safety, tolerability, and immunogenicity of an inactivated SARS-CoV-2 vaccine in healthy adults aged 18–59 years: a randomised, double-blind, placebo-controlled, phase 1/2 clinical trial. *Lancet Infect. Dis.* 21(1):39–51
172. Keech C, Albert G, Cho I, Robertson A, Reed P, et al. 2020. Phase 1–2 trial of a SARS-CoV-2 recombinant spike protein nanoparticle vaccine. *N. Engl. J. Med.* 383:2320–32
173. Graham SP, McLean RK, Spencer AJ, Belij-Rammerstorfer S, Wright D, et al. 2020. Evaluation of the immunogenicity of prime-boost vaccination with the replication-deficient viral vectored COVID-19 vaccine candidate ChAdOx1 nCoV-19. *NPJ Vaccines* 5:69

174. Voysey M, Clemens SAC, Madhi SA, Weckx LY, Folegatti PM, et al. 2020. Safety and efficacy of the ChAdOx1 nCoV-19 vaccine (AZD1222) against SARS-CoV-2: an interim analysis of four randomised controlled trials in Brazil, South Africa, and the UK. *Lancet* 397(10269):99–111
175. Ramasamy MN, Minassian AM, Ewer KJ, Flaxman AL, Folegatti PM, et al. 2020. Safety and immunogenicity of ChAdOx1 nCoV-19 vaccine administered in a prime-boost regimen in young and old adults (COV002): a single-blind, randomised, controlled, phase 2/3 trial. *Lancet* 396:1979–93
176. Tostanoski LH, Wegmann F, Martinot AJ, Loos C, McMahan K, et al. 2020. Ad26 vaccine protects against SARS-CoV-2 severe clinical disease in hamsters. *Nat. Med.* 26:1694–700
177. Bos R, Rutten L, van der Lubbe JEM, Bakkers MJG, Hardenberg G, et al. 2020. Ad26 vector-based COVID-19 vaccine encoding a prefusion-stabilized SARS-CoV-2 Spike immunogen induces potent humoral and cellular immune responses. *NPJ Vaccines* 5:91
178. Zhu FC, Guan XH, Li YH, Huang JY, Jiang T, et al. 2020. Immunogenicity and safety of a recombinant adenovirus type-5-vectored COVID-19 vaccine in healthy adults aged 18 years or older: a randomised, double-blind, placebo-controlled, phase 2 trial. *Lancet* 396:479–88
179. Zhu FC, Li YH, Guan XH, Hou LH, Wang WJ, et al. 2020. Safety, tolerability, and immunogenicity of a recombinant adenovirus type-5 vectored COVID-19 vaccine: A dose-escalation, open-label, non-randomised, first-in-human trial. *Lancet* 395:1845–54
180. Logunov DY, Dolzhikova IV, Zubkova OV, Tukhvatullin AI, Shcheblyakov DV, et al. 2020. Safety and immunogenicity of an rAd26 and rAd5 vector-based heterologous prime-boost COVID-19 vaccine in two formulations: two open, non-randomised phase 1/2 studies from Russia. *Lancet* 396:887–97
181. Xia S, Duan K, Zhang Y, Zhao D, Zhang H, et al. 2020. Effect of an inactivated vaccine against SARS-CoV-2 on safety and immunogenicity outcomes: interim analysis of 2 randomized clinical trials. *JAMA* 324:951–60
182. Wang H, Zhang Y, Huang B, Deng W, Quan Y, et al. 2020. Development of an inactivated vaccine candidate, BBIBP-CorV, with potent protection against SARS-CoV-2. *Cell* 182:713–21 e9
183. Xia S, Zhang Y, Wang Y, Wang H, Yang Y, et al. 2021. Safety and immunogenicity of an inactivated SARS-CoV-2 vaccine, BBIBP-CorV: a randomised, double-blind, placebo-controlled, phase 1/2 trial. *Lancet Infect. Dis.* 21:39–51
184. de Souza N. 2018. Organoids. *Nat. Methods* 15:23
185. Lukassen S, Chua RL, Trefzer T, Kahn NC, Schneider MA, et al. 2020. SARS-CoV-2 receptor ACE 2 and TMPRSS 2 are primarily expressed in bronchial transient secretory cells. *EMBO J.* 39:e105114
186. Blanco-Melo D, Nilsson-Payant BE, Liu WC, Uhl S, Hoagland D, et al. 2020. Imbalanced host response to SARS-CoV-2 drives development of COVID-19. *Cell* 181:1036–45.e9
187. Ma D, Chen CB, Jhanji V, Xu C, Yuan XL, et al. 2020. Expression of SARS-CoV-2 receptor ACE2 and TMPRSS2 in human primary conjunctival and pterygium cell lines and in mouse cornea. *Eye* 34:1212–19
188. Appelberg S, Gupta S, Svensson Akusjärvi S, Ambikan AT, Mikaeloff F, et al. 2020. Dysregulation in Akt/mTOR/HIF-1 signaling identified by proteo-transcriptomics of SARS-CoV-2 infected cells. *Emerg. Microbes Infect.* 9:1748–60
189. Shang J, Wan Y, Luo C, Ye G, Geng Q, et al. 2020. Cell entry mechanisms of SARS-CoV-2. *PNAS* 117:11727–34
190. Xia S, Liu M, Wang C, Xu W, Lan Q, et al. 2020. Inhibition of SARS-CoV-2 (previously 2019-nCoV) infection by a highly potent pan-coronavirus fusion inhibitor targeting its spike protein that harbors a high capacity to mediate membrane fusion. *Cell Res.* 30:343–55
191. Buchrieser J, Dufloo J, Hubert M, Monel B, Planas D, et al. 2020. Syncytia formation by SARS-CoV-2-infected cells. *EMBO J.* 39:e106267
192. Wang C, Li W, Drabek D, Okba NM, van Haperen R, et al. 2020. A human monoclonal antibody blocking SARS-CoV-2 infection. *Nat. Commun.* 11:2251
193. Mirabelli C, Wotring JW, Zhang CJ, McCarty SM, Fursmidt R, et al. 2020. Morphological cell profiling of SARS-CoV-2 infection identifies drug repurposing candidates for COVID-19. *bioRxiv* 2020.05.27.117184. <https://www.biorxiv.org/content/10.1101/2020.05.27.117184v4>
194. Liu Y, Hu G, Wang Y, Zhao X, Ji F, et al. 2020. Functional and genetic analysis of viral receptor ACE2 orthologs reveals a broad potential host range of SARS-CoV-2. *bioRxiv* 2020.04.22.046565. <https://www.biorxiv.org/content/10.1101/2020.04.22.046565v4>

195. Vogel A, Kanevsky I, Che Y, Swanson K, Muik A, et al. 2020. A prefusion SARS-CoV-2 spike RNA vaccine is highly immunogenic and prevents lung infection in non-human primates. *bioRxiv* 2020.09.08.280818. <https://www.biorxiv.org/content/10.1101/2020.09.08.280818v1>
196. Ganneru B, Jogdand H, Dharam VK, Molugu NR, Prasad SD, et al. 2020. Evaluation of safety and immunogenicity of an adjuvanted, TH-1 skewed, whole virion inactivated SARS-CoV-2 vaccine-BBV152. *bioRxiv* 2020.09.09.285445. <https://www.biorxiv.org/content/10.1101/2020.09.09.285445v2>
197. Ella R, Reddy S, Jogdand H, Sarangi V, Ganneru B, et al. 2020. Safety and immunogenicity clinical trial of an inactivated SARS-CoV-2 vaccine, BBV152 (a phase 2, double-blind, randomised controlled trial) and the persistence of immune responses from a phase 1 follow-up report. *medRxiv* 2020.12.21.20248643. <https://www.medrxiv.org/content/10.1101/2020.12.21.20248643v1>
198. Mohandas S, Yadav PD, Shete A, Abraham P, Mohan K, et al. 2020. Immunogenicity and protective efficacy of BBV152: a whole virion inactivated SARS CoV-2 vaccine in the Syrian hamster model. *Res. Sq.* 76768. <https://www.researchsquare.com/article/rs-76768/v1>
199. Yadav P, Ella R, Kumar S, Patil D, Mohandas S, et al. 2020. Remarkable immunogenicity and protective efficacy of BBV152, an inactivated SARS-CoV-2 vaccine in rhesus macaques. *Res. Sq.* 65715. <https://www.researchsquare.com/article/rs-65715/v1>